

PCT

WO. INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



6

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12Q 1/68</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/12760</b> (43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19768 (22) International Filing Date: 27 August 1999 (27.08.99) (30) Priority Data: 09/141,825 28 August 1998 (28.08.98) US 09/172,711 13 October 1998 (13.10.98) US 09/172,108 13 October 1998 (13.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 09/141,825 (CIP) Filed on 28 August 1998 (28.08.98) US 09/172,711 (CIP) Filed on 13 October 1998 (13.10.98) US 09/172,108 (CIP) Filed on 13 October 1998 (13.10.98) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CUNNINGHAM, Mary, Jane [US/US]; 1204 Manet Drive, Sunnyvale, CA 94087			(US). ZWEIGER, Gary, B. [US/US]; 2700 Del Medio Court, #124, Mountain View, CA 95023 (US). PANZER, Scott, R. [US/US]; 965 East El Camino, #621, Sunnyvale, CA 94087 (US). SEILHAMER, Jeffrey, J. [US/US]; 12555 La Cresta, Los Altos, CA 94022 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TOXICOLOGICAL RESPONSE MARKERS			
(57) Abstract <p>The present invention relates to a composition comprising a plurality of nucleic acid molecules. The composition can be used as hybridizable array elements in a microarray. The present invention also relates to methods for screening compounds and therapeutics for metabolic responses indicative of a toxic compound.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## TOXICOLOGICAL RESPONSE MARKERS

This application is filled under the Patent Cooperation Treaty and claims the benefits of U.S. Nonprovisional Application No. 09/141,825, our Docket No. PA-0010 US, filed on August 28, 1998, U.S. Nonprovisional Application No. 09/172,711, our Docket No. PA-0011 US, filed on October 13, 1998, and U.S. Nonprovisional Application No. 09/172,108, our Docket No. PA-0012 US, filed on October 13, 1998.

### TECHNICAL FIELD

The present invention relates to compositions and methods for use in detecting metabolic and toxicological responses.

### BACKGROUND ART

Toxicity testing is a mandatory and time-consuming part of the pharmaceutical drug development pipeline. A more rapid screen to determine the effects upon metabolism and to detect toxicity of lead drug candidates may be the use of gene expression microarrays. For example, microarrays of various kinds may be produced using full length genes or gene fragments. These arrays can then be used to test samples treated with the drug candidates to elucidate the gene expression pattern associated with drug treatment. This gene pattern can be compared with gene expression patterns associated with compounds which produce known toxicological and metabolic responses.

Benzo(a)pyrene is a known rodent and likely human carcinogen and is the prototype of a class of compounds, the polycyclic aromatic hydrocarbons (PAH). It is metabolized by several forms of cytochrome P450 (P450 isozymes) and associated enzymes to form both activated and detoxified metabolites. The ultimate metabolites are the bay-region diol epoxide, benzo(a)pyrene-7,8-diol-9,10-epoxide (BPDE) and the K-region diol epoxide, 9-hydroxy benzo(a)pyrene-4,5-oxide, both of which induce formation of DNA adducts. DNA adducts have been shown to persist in rat liver up to 56 days following treatment with benzo(a)pyrene at a dose of 10 mg/kg body weight three times per week for two weeks (Qu and Stacey (1996) Carcinogenesis 17:53-59).

Acetaminophen is a widely-used analgesic. It is metabolized by specific cytochrome P450 isozymes with the majority of the drug undergoing detoxification by glucuronic acid, sulfate and glutathione conjugation pathways. However, at supratherapeutic doses, acetaminophen is metabolized to an active intermediate, *N*-acetyl-*p*-benzoquinone imine (NAPQI) which can cause hepatic and renal failure. NAPQI then binds to sulfhydryl groups of proteins causing their inactivation and leading to subsequent cell death (Kroger et al. (1997) Gen. Pharmacol. 28:257-263).

Clofibrate is an hypolipidemic drug which lowers elevated levels of serum triglycerides.

In rodents, chronic treatment produces hepatomegaly and an increase in hepatic peroxisomes. Clofibrate has been shown to increase levels of cytochrome P450 4A and reduce the levels of P450 4F. It is also involved in transcription of  $\beta$ -oxidation genes as well as induction of peroxisome proliferator (PP) activated receptors (Kawashima *et al.* (1997) Arch. Biochem. Biophys. 347:148-154). Peroxisome proliferation that is induced by both clofibrate and the chemically-related compound fenofibrate is mediated by a common inhibitory effect on mitochondrial membrane depolarization (Zhou and Wallace (1999) Toxicol. Sci. 48:82-89).

The present invention provides compositions and methods for the screening of compounds for metabolic and toxicological responses.

#### DISCLOSURE OF INVENTION

The invention provides nucleic acid molecules whose transcript levels are modulated in a sample during a metabolic response to a toxic compound. The invention also provides nucleic acid molecules whose transcript levels are upregulated in a sample during a metabolic response to a toxic compound. The invention also provides nucleic acid molecules whose transcript levels are downregulated in a sample during a metabolic response to a toxic compound. Upregulation or downregulation is at least 2 fold, more preferably at least 2.5 fold, most preferably at least 3 fold. The metabolic response to a toxic compound may be a toxicological response.

In another aspect, the invention provides a method for screening a compound for a metabolic response to a test compound or molecule. The method comprises treating a tissue with a known toxic compound, determining levels of a plurality of nucleic acid molecules, selecting from the plurality of nucleic acid molecules those nucleic acid molecules that have levels modulated in samples treated with known toxic compounds when compared with untreated samples. Some of the transcript levels may be upregulated by a toxic compound, others may be downregulated by a toxic compound, and still others may be upregulated with one known toxic compound and be downregulated with another known toxic compound. The selected nucleic acid molecules which are upregulated and downregulated by a known toxic compound are arrayed upon a substrate. The method further comprises determining levels of nucleic acid molecules in a sample after the sample is treated with a compound. Levels of nucleic acid molecules in a sample so treated are then compared with the plurality of the arrayed nucleic acid molecules to identify which sample nucleic acid molecules are upregulated and downregulated by the compound.

Preferably, the comparing comprises contacting the arrayed nucleic acid molecules with the sample nucleic acid molecules under conditions effective to form hybridization complexes between the arrayed nucleic acid molecules and the sample nucleic acid molecules; and detecting the presence or absence of the hybridization complexes. In this context, similarity may mean that at least 1, preferably at least 5, more preferably at least 10, of the upregulated arrayed nucleic acid



molecules form hybridization complexes with the sample nucleic acid molecules at least once during a time course to a greater extent than would the probes derived from a sample not treated with the test compound or a known toxic compound. Similarity may also mean that at least 1, preferably at least 3, of the downregulated arrayed nucleic acid molecules form hybridization  
5 complexes with the sample nucleic acid molecules at least once during a time course to a lesser extent than would the sample nucleic acid molecules of a sample not treated with the test compound or a known toxic compound.

Preferred tissues are selected from the group consisting of liver, kidney, brain, spleen, pancreas and lung. Preferred toxic compounds are selected from the group consisting of  
10 hypolipidemic drugs, n-alkylcarboxylic acids, n-alkylcarboxylic acid precursors, azole antifungal compounds, leukotriene D4 antagonists, herbicides, pesticides, phthalate esters, phenyl acetate, dehydroepiandrosterone sulfate, oleic acid, methanol and their corresponding metabolites, acetaminophen and its corresponding metabolites, benzo(a)pyrene, 3-methylcholanthrene, benz(a)anthracene, 7,12-dimethylbenz(a)anthracene, their corresponding metabolites, and the like.  
15 The arrayed nucleic acid molecules comprise fragments of messenger RNA transcripts of genes that are up-regulated or down-regulated at least 2-fold, preferably at least 2.5-fold, more preferably at least 3-fold, in samples treated with known toxic compounds when compared with untreated samples. Preferred arrayed nucleic acid molecules are selected from the group consisting of SEQ ID NOs:1-117, or fragments thereof, some of whose levels are upregulated and  
20 others of whose levels are downregulated. More preferable are SEQ ID NOs: 3, 9, 10, 13, 19, 26, 31, 33, 35, 36, 37, 39, 42, 57, 67, 78, 81, 82, 94, and 98 which are upregulated, and SEQ ID NOs: 43, 49, 50, 52, 53, 54, 55, 56, 59, 61, 63, 68, 71, 74, 85, 87, 90, 95, 102, 103, 105, and 115 which are downregulated. Most preferable are SEQ ID NOs: 31, 33, 35, 36, 39, 52, 53, 54, 55, 63, 74, 81, 90, 94, and 95. In one embodiment, the polynucleotide targets are hybridizable array elements  
25 of a microarray.

Alternatively, the invention provides methods for screening a test compound or molecule for a metabolic response or for screening a sample for a metabolic response to a test compound or molecule.

Alternatively, the invention provides methods for screening a test compound or molecule  
30 for a previously unknown metabolic response or for screening a sample for a previously unknown metabolic response to a test compound or molecule.

In another aspect, the invention provides methods for preventing a toxicological response by administering complementary nucleotide molecules against one or more selected upregulated nucleic acid molecules or a ribozyme that specifically cleaves such molecules. Alternatively, a  
35 toxicological response may be prevented by administering sense nucleotide molecules for one or

more selected downregulated nucleic acid molecules.

In yet another aspect, the invention provides methods for preventing a toxicological response by administering an agonist which initiates transcription of a gene comprising a downregulated nucleic acid molecule of the invention. Alternatively, a toxicological response may  
5 be prevented by administering an antagonist which prevents transcription of a gene comprising an upregulated nucleic acid molecule of the invention.

The invention also provides a substantially purified mammalian protein or a portion thereof. The invention further provides isolated and purified proteins encoded by the nucleic acid molecules of SEQ ID NOs:1-117. Additionally, the invention provides a pharmaceutical  
10 composition comprising a substantially purified mammalian protein or a portion thereof in conjunction with a pharmaceutical carrier.

The invention further provides a method for using at least a portion of the proteins encoded by SEQ ID NOs:1-117 to produce antibodies. The invention also provides a method for using a protein or a portion thereof to screen a library of molecules to identify at least one ligand  
15 which specifically binds the protein, the method comprising combining the protein with the library of molecules under conditions allowing specific binding, and detecting specific binding, thereby identifying a ligand which specifically binds the protein. Such libraries include DNA and RNA molecules, peptides, agonists, antagonists, antibodies, immunoglobulins, drug compounds, pharmaceutical agents, and other ligands. In one aspect, the ligand identified using the method  
20 modulates the activity of the mammalian protein. In an analogous method, the protein or a portion thereof is used to purify a ligand. The method involves combining the protein or a portion thereof with a sample under conditions to allow specific binding, detecting specific binding between the protein and ligand, recovering the bound protein, and separating the protein from the ligand to obtain purified ligand.

The invention further provides a method for inserting a marker gene into the genomic DNA of an animal to disrupt the expression of the natural nucleic acid molecule. The invention also provides a method for using the nucleic acid molecule to produce an animal model system, the method comprising constructing a vector containing the nucleic acid molecule; introducing the vector into a totipotent embryonic stem cell; selecting an embryonic stem cell with the vector  
30 integrated into genomic DNA; microinjecting the selected cell into a blastocyst, thereby forming a chimeric blastocyst; transferring the chimeric blastocyst into a pseudopregnant dam, wherein the dam gives birth to a chimeric animal containing at least one additional copy of nucleic acid molecule in its germ line; and breeding the chimeric animal to generate a homozygous animal model system.

35

### BRIEF DESCRIPTION OF THE SEQUENCE LISTING

A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure, as it appears in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever.

The Sequence Listing contains the nucleic acid sequence of exemplary nucleic acid molecules of the invention, SEQ ID NOs: 1-117.

### MODES FOR CARRYING OUT THE INVENTION

#### 10 Definitions

"Sample" is used in its broadest sense. A sample containing nucleic acid molecules may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; genomic DNA, RNA, or cDNA in solution or bound to a substrate; a cell; a biological tissue or isolated fragment thereof, for example, a needle biopsy; a fingerprint or tissue print; natural or synthetic fibres; in a solution; in a liquid suspension; in a gaseous suspension; in an aerosol; and the like.

"Plurality" refers preferably to a group of one or more members, preferably to a group of at least about 10, and more preferably to a group of at least about 100 members, and even more preferably a group of 10,000 members.

20 "Substrate" refers to a rigid or semi-rigid support to which nucleic acid molecules or proteins are bound and includes membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, capillaries or other tubing, plates, polymers, and microparticles with a variety of surface forms including wells, trenches, pins, channels and pores.

"Modulates" refers to a change in activity (biological, chemical, or immunological) or lifespan resulting from specific binding between a molecule and either a nucleic acid molecule or a protein

"Microarray" refers to an ordered arrangement of hybridizable array elements on a substrate. The array elements are arranged so that there are preferably at least ten or more different array elements, more preferably at least 100 array elements, even more preferably at least 1000 array elements, and most preferably 10,000. Furthermore, the hybridization signal from each of the array elements is individually distinguishable. In a preferred embodiment, the array elements comprise nucleic acid molecules.

"Nucleic acid molecule" refers to a nucleic acid, oligonucleotide, nucleotide, polynucleotide or any fragment thereof. It may be DNA or RNA of genomic or synthetic origin, double-stranded or single-stranded, and combined with carbohydrate, lipids, protein or other

materials to perform a particular activity such as transformation or form a useful composition such as a peptide nucleic acid (PNA). "Oligonucleotide" is substantially equivalent to the terms amplimer, primer, oligomer, element, target, and probe and is preferably single stranded.

"Protein" refers to an amino acid sequence, oligopeptide, peptide, polypeptide or portions thereof whether naturally occurring or synthetic.

"Up-regulated" refers to a nucleic acid molecule whose levels increased in a treated sample compared with the nucleic acid molecule in an untreated sample.

"Down-regulated" refers to nucleic acid molecule whose levels decreased in a treated sample compared with the nucleic acid molecule in an untreated sample.

"Toxic compound" or "toxic agent" is any compound, molecule, or agent that elicits a biochemical, metabolic, and physiological response in an individual or animal, such as i) DNA damage, ii) cell damage, iii) organ damage or cell death, or iv) clinical morbidity or mortality.

"Toxicological response" refers to a biochemical, metabolic, and physiological response in an individual, animal, or test system which has been exposed to a toxic compound or toxic agent.

"Fragment" refers to an Incyte clone or any part of a nucleic acid molecule which retains a usable, functional characteristic. Useful fragments include oligonucleotides and polynucleotides which may be used in hybridization or amplification technologies or in regulation of replication, transcription or translation. Exemplary fragments are the first twenty consecutive nucleotides of SEQ ID NOs:1-117.

"Hybridization complex" refers to a complex between two nucleic acid molecules by virtue of the formation of hydrogen bonds between purines and pyrimidines.

"Ligand" refers to any molecule, agent, or compound which will bind specifically to a complementary site on a nucleic acid molecule or protein. Such ligands stabilize or modulate the activity of nucleic acid molecules or proteins of the invention and may be composed of at least one of the following: inorganic and organic substances including nucleic acids, proteins, carbohydrates, fats, and lipids.

"Substantially purified" refers to nucleic acid molecules or proteins that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free, from other components with which they are naturally associated.

### **The Invention**

The present invention provides a composition and method of using the composition for screening test compounds and molecules for toxicological responses. Additionally the invention provides methods for characterizing the toxicological responses of a sample to a test compound or molecule. In particular, the present invention provides a composition comprising a plurality of

nucleic acid molecules derived from human cDNA libraries, monkey cDNA libraries, mouse cDNA libraries, normal rat liver cDNA libraries, normalized rat liver cDNA libraries and prehybridized rat liver cDNA libraries and rat kidney cDNA libraries in a test system. The nucleic acid molecules have been further selected for exhibiting up-regulated or down-regulated gene expression in rat livers when the rats have been exposed to a known hepatotoxin, including a peroxisome proliferator (PP), acetaminophen or one of its corresponding metabolites, and a polycyclic aromatic hydrocarbon (PAH).

PPs include hypolipidemic drugs, such as clofibrate, fenofibrate, clofenic acid, nafenopin, gemfibrozil, ciprofibrate, bezafibrate, halofenate, simfibrate, benzofibrate, etofibrate, WY-14,643, and the like; n-alkylcarboxylic acids, such as trichloroacetic acid, valproic acid, hexanoic acid, and the like; n-alkylcarboxylic acid precursors, such as trichloroethylene, etrachloroethylene, and the like; azole antifungal compounds, such as bifenazole, and the like; leukotriene D4 antagonists; herbicides; pesticides; phthalate esters, such as di-[2-ethylhexyl] phthalate, mono-[2-ethylhexyl] phthalate, and the like; and natural chemicals, such as phenyl acetate, dehydroepiandrosterone sulfate, oleic acid, methanol, and the like. In a preferred embodiment the toxic compound is clofibrate, or one of its corresponding metabolites. In another preferred embodiment the toxic compound is fenofibrate, or one of its corresponding metabolites.

PAHs include compounds such as benzo(a)pyrene, 3-methylcholanthrene, benzo(a)anthracene, 7,12-dimethylbenz(a)anthracene, their corresponding metabolites, and the like. In a preferred embodiment the toxic compound is benzo(a)pyrene, or one of its corresponding metabolites.

SEQ ID NOs:1-117 were identified by their pattern of at least two-fold up-regulation or down-regulation following hybridization with sample nucleic acid molecules from treated rat liver tissue. These and other nucleic acid molecules can be immobilized on a substrate as hybridizable array elements in a microarray format. The microarray may be used to characterize gene expression patterns associated with novel compounds to elucidate any metabolic responses or to monitor the effects of treatments during clinical therapy where metabolic responses to toxic compounds may be expected.

When the nucleic acid molecules are employed as hybridizable array elements in a microarray, the array elements are organized in an ordered fashion so that each element is present at a specified location on the substrate. Because the array elements are at specified locations on the substrate, the hybridization patterns and intensities (which together create a unique expression profile) can be interpreted in terms of expression levels of particular genes and can be correlated with a toxicological response associated with a test compound or molecule.

Furthermore, the present invention provides methods for screening test compounds and/or

molecules for potential toxicological responses and for screening a sample's toxicological response to a particular test compound or molecule. Briefly, these methods entail treating a sample with the test compound or molecule to elicit a change in gene expression patterns comprising the expression of a plurality of sample nucleic acid molecules. Nucleic acid molecules are selected by identifying those levels of expressed nucleic acid molecules in rat liver or kidney which are up-regulated or down-regulated at least 2-fold, more preferably at least 2.5-fold, most preferably at least 3-fold, when treated with a known toxic compound. The nucleic acid molecules are arrayed on a substrate. Then, the arrayed nucleic acid molecules and sample nucleic acid molecules are combined under conditions effective to form hybridization complexes which may be detected by methods well known in the art. Detection of higher or lower levels of such hybridization complexes compared with hybridization complexes derived from samples treated with a compound that is known not to induce a toxicological response correlates with a toxicological response to a test compound or a toxicological response to a molecule.

#### Complementary DNA libraries

Molecules are identified that reflect all or most of the genes that are expressed in rat tissue. Molecules may be identified by isolating clones derived from several types of rat cDNA libraries, including normal rat cDNA libraries, normalized rat cDNA libraries and prehybridized rat cDNA libraries. Clone inserts derived from these clones may be partially sequenced to generate expressed sequence tags (ESTs).

In one embodiment, two collections of ESTs are identified and sequenced. A first collection of ESTs (the originator molecules) are derived from rat liver and kidney and from the cDNA libraries presented in the Examples. A second collection includes ESTs derived from other rat cDNA libraries available in the ZOOSEQ database (Incyte Pharmaceuticals, Inc., Palo Alto CA).

The two collections of ESTs are clustered electronically to form master clusters of ESTs. Master clusters are formed by identifying overlapping EST molecules and assembling these ESTs. A nucleic acid fragment assembly tool, such as the Phrap tool (Phil Green, University of Washington) and the GELVIEW fragment assembly system (GCG, Madison WI), can be used for this purpose. The minimum number of clones which constitute a cluster is two. In another embodiment, a collection of human genes known to be expressed in response to toxic agents are used to select representative ESTs from the 113 rat cDNA libraries. The master cluster process is repeated for these molecules.

After assembling the clustered consensus nucleic acid sequences, a representative 5' clone is nominated from each master cluster. The most 5' clone is preferred because it is most likely to contain the complete gene. The nomination process is described in greater detail in "Relational

Database and System for Storing Information Relating to Biomolecular Sequences and Reagents", USSN 09/034,807, filed March 4, 1998, herein incorporated in its entirety by reference. The EST molecules are used as array elements on a microarray.

Selection of arrayed nucleic acid molecules

5           Samples are treated, preferably at subchronic doses, with one or more known toxic compounds over a defined time course. Preferably, the agents are peroxisomal proliferators (PPs), acetaminophen or one of its corresponding metabolites, and polycyclic aromatic hydrocarbons (PAHs).

10           The gene expression patterns derived from such treated biological samples can be compared with the gene expression patterns derived from untreated biological samples to identify nucleic acid molecules whose expression is either up-regulated or down-regulated due to the response to the toxic compounds. These molecules may then be employed as array elements alone or in combination with other array element molecules. Such a microarray is particularly useful to detect and characterize gene expression patterns associated with known toxic compounds. Such  
15           gene expression patterns can then be used for comparison to identify other compounds which also elicit a metabolic response to a toxic compound.

          The arrayed nucleic acid molecules can be manipulated to optimize their performance in hybridization. To optimize hybridization, the arrayed nucleic acid molecules are examined using a computer algorithm to identify portions of genes without potential secondary structure. Such  
20           computer algorithms are well known in the art and are part of OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or LASERGENE software (DNASTAR, Madison WI). These programs can search within nucleic acid molecule sequences to identify stem loop structures and tandem repeats and to analyze G + C content of the sequence (those molecules with a G + C content greater than 60% are excluded). Alternatively, the arrayed nucleic acid molecules  
25           can be optimized by trial and error. Experiments can be performed to determine whether sample nucleic acid molecules and complementary arrayed nucleic acid molecules hybridize optimally under experimental conditions.

          The arrayed nucleic acid molecules can be any RNA-like or DNA-like material, such as mRNAs, cDNAs, genomic DNA, peptide nucleic acids, branched DNAs and the like. The arrayed  
30           nucleic acid molecules can be in sense or antisense orientations.

          In one embodiment, the arrayed nucleic acid molecules are cDNAs. The size of the DNA sequence of interest may vary, and is preferably from 50 to 10,000 nucleotides, more preferably from 150 to 3,500 nucleotides. In a second embodiment, the nucleic acid molecules are vector DNAs. In this case the size of the DNA sequence of interest, i.e., the insert sequence, may vary  
35           from about 50 to 10,000 nucleotides, more preferably from about 150 to 3,500 nucleotides.

The nucleic acid molecule sequences of the Sequence Listing have been prepared by current, state-of-the-art, automated methods and, as such, may contain occasional sequencing errors and unidentified nucleotides. Nucleotide analogues can be incorporated into the nucleic acid molecules by methods well known in the art. The only requirement is that the incorporated nucleotide analogues must serve to base pair with sample nucleic acid molecules. For example, certain guanine nucleotides can be substituted with hypoxanthine which base pairs with cytosine residues. However, these base pairs are less stable than those between guanine and cytosine. Alternatively, adenine nucleotides can be substituted with 2,6-diaminopurine which can form stronger base pairs than those between adenine and thymidine. Additionally, the nucleic acid molecules can include nucleotides that have been derivatized chemically or enzymatically. Typical modifications include derivatization with acyl, alkyl, aryl or amino groups.

The nucleic acid molecules can be immobilized on a substrate via chemical bonding. Furthermore, the molecules do not have to be directly bound to the substrate, but rather can be bound to the substrate through a linker group. The linker groups are typically about 6 to 50 atoms long to provide exposure to the bound nucleic acid molecule. Preferred linker groups include ethylene glycol oligomers, diamines, diacids and the like. Reactive groups on the substrate surface react with one of the terminal portions of the linker to bind the linker to the substrate. The other terminal portion of the linker is then functionalized for binding the nucleic acid molecule. Preferred substrates are any suitable rigid or semirigid support, including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which the arrayed nucleic acid molecules are bound.

The samples can be any sample comprising sample nucleic acid molecules and obtained from any bodily fluid (blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. The samples can be derived from any species, but preferably from eukaryotic species, and more preferably from mammalian species such as rat and human.

DNA or RNA can be isolated from the sample according to any of a number of methods well known to those of skill in the art. For example, methods of purification of nucleic acids are described in Laboratory Techniques in Biochemistry and Molecular Biology: Hybridization With Nucleic Acid Probes, Part I. Theory and Nucleic Acid Preparation, P. Tijssen, ed. Elsevier (1993). In one preferred embodiment, total RNA is isolated using the TRIZOL total RNA isolation reagent (Life Technologies, Gaithersburg MD) and mRNA is isolated using oligo d(T) column chromatography or glass beads. When sample nucleic acid molecules are amplified it is desirable to amplify the sample nucleic acid molecules and maintain the relative abundances of the original sample, including low abundance transcripts. RNA can be amplified in vitro, in situ or in vivo.



(See Eberwine US Patent No. 5,514,545).

It is also advantageous to include controls within the sample to assure that amplification and labeling procedures do not change the true distribution of nucleic acid molecules in the sample. For this purpose, a sample is spiked with an amount of control nucleic acid molecules  
5 predetermined to be detectable upon hybridization to its complementary arrayed nucleic acid molecule and the composition of nucleic acid molecules includes reference nucleic acid molecules which specifically hybridize with the control arrayed nucleic acid molecules. After hybridization and processing, the hybridization signals obtained should reflect accurately the amounts of control arrayed nucleic acid molecules added to the sample.

10 Prior to hybridization, it may be desirable to fragment the sample nucleic acid molecules. Fragmentation improves hybridization by minimizing secondary structure and cross-hybridization to other sample nucleic acid molecules in the sample or noncomplementary nucleic acid molecules. Fragmentation can be performed by mechanical or chemical means.

#### Labeling

15 The sample nucleic acid molecules may be labeled with one or more labeling moieties to allow for detection of hybridized arrayed/sample nucleic acid molecule complexes. The labeling moieties can include compositions that can be detected by spectroscopic, photochemical, biochemical, bioelectronic, immunochemical, electrical, optical or chemical means. The labeling moieties include radioisotopes, such as  $^{32}\text{P}$ ,  $^{33}\text{P}$  or  $^{35}\text{S}$ , chemiluminescent compounds, labeled  
20 binding proteins, heavy metal atoms, spectroscopic markers, such as fluorescent markers and dyes, magnetic labels, linked enzymes, mass spectrometry tags, spin labels, electron transfer donors and acceptors, and the like. Preferred fluorescent markers include Cy3 and Cy5 fluorophores (Amersham Pharmacia Biotech, Piscataway NJ).

#### Hybridization

25 The nucleic acid molecule sequence of SEQ ID NOs:1-117 and fragments thereof can be used in various hybridization technologies for various purposes in a test system. Hybridization probes may be designed or derived from SEQ ID NOs:1-117. Such probes may be made from a highly specific region such as the 5' regulatory region or from a conserved motif, and used in protocols to identify naturally occurring sequences encoding the mammalian protein, allelic  
30 variants, or related sequences, and should preferably have at least 50% sequence identity to any of the protein sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NOs:1-117 or from genomic sequences including promoters, enhancers, and introns of the mammalian gene. Hybridization or PCR probes may be produced using oligolabeling, nick translation, end-labeling, or PCR amplification in the  
35 presence of the labeled nucleotide. A vector containing the nucleic acid sequence may be used to

produce an mRNA probe in vitro by addition of an RNA polymerase and labeled nucleic acid molecules. These procedures may be conducted using commercially available kits such as those provided by Amersham Pharmacia Biotech.

The stringency of hybridization is determined by G+C content of the probe, salt concentration, and temperature. In particular, stringency can be increased by reducing the concentration of salt or raising the hybridization temperature. In solutions used for some membrane based hybridizations, addition of an organic solvent such as formamide allows the reaction to occur at a lower temperature. Hybridization can be performed at low stringency with buffers, such as 5 x SSC with 1% sodium dodecyl sulfate (SDS) at 60°C, which permits the formation of a hybridization complex between nucleotide sequences that contain some mismatches. Subsequent washes are performed at higher stringency with buffers such as 0.2 x SSC with 0.1% SDS at either 45°C (medium stringency) or 68°C (high stringency). At high stringency, hybridization complexes will remain stable only where the nucleic acid sequences are completely complementary. In some membrane-based hybridizations, preferably 35% or most preferably 50%, formamide can be added to the hybridization solution to reduce the temperature at which hybridization is performed, and background signals can be reduced by the use of other detergents such as Sarkosyl or Triton X-100 and a blocking agent such as salmon sperm DNA. Selection of components and conditions for hybridization are well known to those skilled in the art and are reviewed in Ausubel (supra) and Sambrook et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY.

Hybridization specificity can be evaluated by comparing the hybridization of specificity-control nucleic acid molecules to specificity-control sample nucleic acid molecules that are added to a sample in a known amount. The specificity-control arrayed nucleic acid molecules may have one or more sequence mismatches compared with the corresponding arrayed nucleic acid molecules. In this manner, whether only complementary arrayed nucleic acid molecules are hybridizing to the sample nucleic acid molecules or whether mismatched hybrid duplexes are forming is determined.

Hybridization reactions can be performed in absolute or differential hybridization formats. In the absolute hybridization format, nucleic acid molecules from one sample are hybridized to the molecules in a microarray format and the signals detected after hybridization complex formation correlate to nucleic acid molecule levels in a sample. In the differential hybridization format, the differential expression of a set of genes in two biological samples is analyzed. For differential hybridization, nucleic acid molecules from both biological samples are prepared and labeled with different labeling moieties. A mixture of the two labeled nucleic acid molecules is added to a microarray. The microarray is then examined under conditions in which the emissions from the

two different labels are individually detectable. Molecules in the microarray that are hybridized to substantially equal numbers of nucleic acid molecules derived from both biological samples give a distinct combined fluorescence (Shalon *et al.* PCT publication WO95/35505). In a preferred embodiment, the labels are fluorescent markers with distinguishable excitation and emission spectra, such as Cy3 and Cy5 fluorophores.

After hybridization, the microarray is washed to remove nonhybridized nucleic acid molecules, then complex formation between the hybridizable array elements and the nucleic acid molecules is detected. Methods for detecting complex formation are well known to those skilled in the art. In a preferred embodiment, the nucleic acid molecules are labeled with a fluorescent label and measurement of levels and patterns of fluorescence indicative of complex formation is accomplished by fluorescence microscopy, preferably confocal fluorescence microscopy.

In a differential hybridization experiment, nucleic acid molecules from two or more different biological samples are labeled with two or more different fluorescent labels with different excitation and emission wavelengths. The labeled sample is excited with a specific excitation wavelength. Fluorescent signals are detected separately with different photomultipliers set to detect specific emission wavelengths. The relative abundances/expression levels of the nucleic acid molecules in two or more samples is obtained.

Typically, microarray fluorescence intensities can be normalized to take into account variations in hybridization intensities when more than one microarray is used under similar test conditions. In a preferred embodiment, individual arrayed-sample nucleic acid molecule complex hybridization intensities are normalized using the intensities derived from internal normalization controls contained on each microarray.

The labeled sample emits specific wavelengths which are detected using a plurality of photomultipliers. The relative abundances/expression levels of the arrayed nucleic acid molecules can be used as hybridizable elements in a microarray. Such a microarray can be employed to identify expression profiles associated with particular toxicological responses. Then, a particular subset of these photomultipliers are set to detect specific wavelengths. The relative expression levels of the arrayed nucleic acid molecules can be identified as to which arrayed nucleic acid molecule expression is modulated in response to a particular toxicological agent. These photomultipliers set to detect specific wavelengths. The relative expression levels of the nucleic acid molecules can be employed to identify other compounds with a similar toxicological response.

Alternatively, for some treatments with known side effects, the microarray, and expression patterns derived therefrom, is employed to "fine tune" the treatment regimen. A dosage is established that minimizes expression patterns associated with undesirable side effects. This

approach may be more sensitive and rapid than waiting for the patient to show toxicological side effects before altering the course of treatment.

Generally, the method for screening a library of test compounds or molecules to identify those with a toxicological response entails selecting a plurality of arrayed nucleic acid molecule  
5 genes whose expression levels are modulated in tissues treated with known toxic compounds when compared with untreated tissues. Then a sample is treated with the test compound or molecule to induce a pattern of gene expression comprising the expression of a plurality of nucleic acid molecules. A test compound may be screened at several doses to determine which doses may be toxic and which may not.

10 Then, the expression levels of the arrayed nucleic acid molecules and the sample nucleic acid molecules are compared to identify those compounds that induce expression levels of the sample nucleic acid molecules that are similar to those of the arrayed nucleic acid molecules. In one preferred embodiment, gene expression levels are compared by contacting the arrayed nucleic acid molecules with the sample nucleic acid molecules under conditions effective to form  
15 hybridization complexes between arrayed nucleic acid molecules and sample nucleic acid molecules, and detecting the presence or absence of the hybridization complexes.

Similarity may mean that at least 1, preferably at least 5, more preferably at least 10, of the upregulated arrayed nucleic acid molecules form hybridization complexes with the sample nucleic acid molecules at least once during a time course to a greater extent than would the nucleic acid  
20 molecules of a sample not treated with the test compound. Similarity may also mean that at least 1, preferably at least 3, of the downregulated nucleic acid molecules form hybridization complexes with the nucleic acid molecules at least once during a time course to a lesser extent than would the nucleic acid molecules of a sample not treated with the test compound.

Such a similarity of expression patterns means that a toxicological response is associated  
25 with the test compound or molecule tested. Preferably, the toxic compounds belong to the class of peroxisomal proliferators (PPs), including hypolipidemic drugs, such as clofibrate, fenofibrate, clofenic acid, nafenopin, gemfibrozil, ciprofibrate, bezafibrate, halofenate, simfibrate, benzofibrate, etofibrate, WY-14,643, and the like; n-alkylcarboxylic acids, such as trichloroacetic acid, valproic acid, hexanoic acid, and the like; n-alkylcarboxylic acid precursors, such as  
30 trichloroethylene, etrachloroethylene, and the like; azole antifungal compounds, such as bifentazole, and the like; leukotriene D4 antagonists; herbicides; pesticides; phthalate esters, such as di-[2-ethylhexyl] phthalate, mono-[2-ethylhexyl] phthalate, and the like; and natural chemicals, such as phenyl acetate, dehydroepiandrosterone sulfate, oleic acid, methanol, and the like. In another embodiment, the toxic compounds are acetaminophen or one of its corresponding  
35 metabolites. In yet another embodiment, the toxic compound is a polycyclic aromatic hydrocarbon

(PAH), including compounds such as benzo(a)pyrene, 3-methylcholanthrene, benz(a)anthracene, 7,12-dimethylbenz(a)anthracene, their corresponding metabolites, and the like. Of particular interest is the study of the metabolic responses of these compounds on the liver, kidney, brain, spleen, pancreas, and lung.

5    Modification of Gene Expression Using Nucleic Acids

Gene expression may be modified by designing complementary or antisense molecules (DNA, RNA, or PNA) to the control, 5', 3', or other regulatory regions of the mammalian gene. Oligonucleotides designed with reference to the transcription initiation site are preferred. Similarly, inhibition can be achieved using triple helix base-pairing, which inhibits the binding of  
10    polymerases, transcription factors, or regulatory molecules (Gee *et al.* In: Huber and Carr (1994) Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177). A complementary molecule may also be designed to block translation by preventing binding between ribosomes and mRNA. In one alternative, a library of nucleic acid molecules or fragments thereof may be screened to identify those which specifically bind a regulatory, nontranslated sequence.

15       Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA followed by endonucleolytic cleavage at sites such as GUA, GUU, and GUC. Once such sites are identified, an oligonucleotide with the same sequence may be evaluated for secondary structural features which would render the  
20    oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing their hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary nucleic acids and ribozymes of the invention may be prepared via recombinant expression, in vitro or in vivo, or using solid phase phosphoramidite chemical synthesis. In addition, RNA molecules may be modified to increase intracellular stability and  
25    half-life by addition of flanking sequences at the 5' and/or 3' ends of the molecule or by the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. Modification is inherent in the production of PNAs and can be extended to other nucleic acid molecules. The inclusion of nontraditional bases such as inosine, queosine, and wybutosine, or the modification of adenine, cytidine, guanine, thymine, and uridine with acetyl-,  
30    methyl-, thio- groups, renders the molecule less available as a substrate to endogenous endonucleases.

Screening Assays

The nucleic acid molecule encoding the mammalian protein may be used to screen a library of molecules for specific binding affinity. The libraries may be DNA molecules, RNA  
35    molecules, PNAs, peptides, proteins such as transcription factors, enhancers, repressors, and other

ligands which regulate the activity, replication, transcription, or translation of the nucleic acid molecule in the biological system. The assay involves combining the mammalian nucleic acid molecule or a fragment thereof with the library of molecules under conditions allowing specific binding, and detecting specific binding to identify at least one molecule which specifically binds the nucleic acid molecule.

Similarly the mammalian protein or a portion thereof may be used to screen libraries of molecules in any of a variety of screening assays. The portion of the protein employed in such screening may be free in solution, affixed to an abiotic or biotic substrate, or located intracellularly. Specific binding between the protein and molecule may be measured. Depending on the kind of library being screened, the assay may be used to identify DNA, RNA, or PNA molecules, agonists, antagonists, antibodies, immunoglobulins, inhibitors, peptides, proteins, drugs, or any other ligand, which specifically binds the protein. One method for high throughput screening using very small assay volumes and very small amounts of test compound is described in USPN 5,876,946, incorporated herein by reference, which screens large numbers of molecules for enzyme inhibition or receptor binding.

#### Purification of Ligand

The nucleic acid molecule or a fragment thereof may be used to purify a ligand from a sample. A method for using a mammalian nucleic acid molecule or a fragment thereof to purify a ligand would involve combining the nucleic acid molecule or a fragment thereof with a sample under conditions to allow specific binding, detecting specific binding, recovering the bound protein, and using an appropriate agent to separate the nucleic acid molecule from the purified ligand.

Similarly, the protein or a portion thereof may be used to purify a ligand from a sample. A method for using a mammalian protein or a portion thereof to purify a ligand would involve combining the protein or a portion thereof with a sample under conditions to allow specific binding, detecting specific binding between the protein and ligand, recovering the bound protein, and using an appropriate chaotropic agent to separate the protein from the purified ligand.

#### Pharmacology

Pharmaceutical compositions are those substances wherein the active ingredients are contained in an effective amount to achieve a desired and intended purpose. The determination of an effective dose is well within the capability of those skilled in the art. For any compound, the therapeutically effective dose may be estimated initially either in cell culture assays or in animal models. The animal model is also used to achieve a desirable concentration range and route of administration. Such information may then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of protein or inhibitor which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity of such agents may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it may be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indexes are preferred. The data obtained from cell culture assays and animal studies are used in formulating a range of dosage for human use.

### Model Systems

Animal models may be used as bioassays where they exhibit a toxic response similar to that of humans and where exposure conditions are relevant to human exposures. Mammals are the most common models, and most toxicity studies are performed on rodents such as rats or mice because of low cost, availability, and abundant reference toxicology. Inbred rodent strains provide a convenient model for investigation of the physiological consequences of under- or over-expression of genes of interest and for the development of methods for diagnosis and treatment of diseases. A mammal inbred to over-express a particular gene, so that the protein is secreted in milk, may also serve as a convenient source of the protein expressed by that gene.

### Toxicology

Toxicology is the study of the effects of test compounds, molecules, or toxic agents on living systems to identify adverse effects. The majority of toxicity studies are performed on rats or mice to help predict whether adverse effects of agents will occur in humans. Observation of qualitative and quantitative changes in physiology, behavior, homeostatic, developmental, and reproductive processes, and lethality are used to generate profiles of safe or toxic responses and to assess the consequences on human health following exposure to the agent.

Genetic toxicology identifies and analyzes the ability of an agent to produce damage at a cellular or subcellular level. Such genotoxic agents usually have common chemical or physical properties that facilitate interaction with nucleic acids and are most harmful when mutated chromosomes are passed along to progeny. Toxicological studies may identify agents that increase the frequency of structural or functional abnormalities in progeny if administered to either parent before conception, to the mother during pregnancy, or to the developing organism. Mice and rats are most frequently used in these tests because of their short reproductive cycle which produces the number of organisms needed to satisfy statistical requirements.

Acute toxicity tests are based on a single administration of the agent to the subject to determine the symptomology or lethality of the agent. Three experiments are conducted: 1) an initial dose-range-finding experiment, 2) an experiment to narrow the range of effective doses,

and 3) a final experiment for establishing the dose-response curve.

Prolonged toxicity tests are based on the repeated administration of the agent. Rat and dog are commonly used in these studies to provide data from species in different taxonomic orders.

With the exception of carcinogenesis, there is considerable evidence that daily administration of an agent at high-dose concentrations for periods of three to four months will reveal most forms of toxicity in adult animals.

Chronic toxicity tests, with a duration of a year or more, are used to demonstrate either the absence of toxicity or the carcinogenic potential of an agent. When studies are conducted on rats, a minimum of at least one test group plus one control group are used. Animals are quarantined, examined for health, and monitored at the outset and at intervals throughout the experiment.

#### Transgenic Animal Models

Transgenic rodents which overexpress or underexpress a gene of interest may be inbred and used to model human diseases or to test compounds and molecules for therapeutic or toxicological effects. (See USPN 4,736,866; USPN 5,175,383; and USPN 5,767,337; incorporated herein by reference). In some cases, the introduced gene may be activated at a specific time in a specific tissue type during fetal development or postnatally. Expression of the transgene is monitored by analysis of phenotype or tissue-specific mRNA expression, in transgenic animals before, during, and after being challenged with experimental drug therapies.

#### Embryonic Stem Cells

Embryonic stem cells (ES) isolated from rodent embryos retain the potential to form an embryo. When ES cells are placed inside a carrier embryo, they resume normal development and contribute to all tissues of the live-born animal. ES cells are the preferred cells used in the creation of experimental knockout and knockin rodent strains. Mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and are grown under culture conditions well known in the art. Vectors for knockout strains contain a disease gene candidate modified to include a marker gene which disrupts transcription and/or translation *in vivo*. The vector is introduced into ES cells by transformation methods such as electroporation, liposome delivery, microinjection, and the like which are well known in the art. The endogenous rodent gene is replaced by the disrupted disease gene through homologous recombination and integration during cell division. Then transformed ES cells are selected under conditions, identified, and preferably microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains.

ES cells are also used to study the differentiation of various cell types and tissues *in vitro*, such as neural cells, hematopoietic lineages, and cardiomyocytes (Bain *et al.* (1995) Dev. Biol.



168:342-357; Wiles and Keller (1991) Development 111:259-267; and Klug *et al.* (1996) J. Clin. Invest. 98:216-224). Recent developments demonstrate that ES cells derived from human blastocysts may also be manipulated *in vitro* to differentiate into eight separate cell lineages, including endoderm, mesoderm, and ectodermal cell types (Thomson (1998) Science 282:1145-1147).

#### Knockout Analysis

In gene knockout analysis, a region of a human disease gene candidate is enzymatically modified to include a non-mammalian gene such as the neomycin phosphotransferase gene (*neo*; Capecchi (1989) Science 244:1288-1292). The inserted coding sequence disrupts transcription and translation of the targeted gene and prevents biochemical synthesis of the disease candidate protein. The modified gene is transformed into cultured embryonic stem cells (described above), the transformed cells are injected into rodent blastulae, and the blastulae are implanted into pseudopregnant dams. Transgenic progeny are crossbred to obtain homozygous inbred lines.

#### Knockin Analysis

Totipotent ES cells, present in the early stages of embryonic development, can be used to create knockin humanized animals (pigs) or transgenic animal models (mice or rats) of human diseases. With knockin technology, a region of a human gene is injected into animal ES cells, and the human sequence integrates into the animal cell genome by recombination. Totipotent ES cells which contain the integrated human gene are handled as described above. Inbred animals are studied and treated to obtain information on the analogous human condition. These methods have been used to model several human diseases. (See, e.g., Lee *et al.* (1998) Proc. Natl. Acad. Sci. 95:11371-11376; Baudoin *et al.* (1998) Genes Dev. 12:1202-1216; and Zhuang *et al.* (1998) Mol. Cell Biol. 18:3340-3349).

#### Non-Human Primate Model

The field of animal testing deals with data and methodology from basic sciences such as physiology, genetics, chemistry, pharmacology and statistics. These data are paramount in evaluating the effects of test compounds or molecules on non-human primates as they can be related to human health. Monkeys are used as human surrogates in vaccine and drug evaluations, and their responses are relevant to human exposures under similar conditions. *Cynomolgus* and rhesus monkeys (*Macaca fascicularis* and *Macaca mulatta*, respectively) and common marmosets (*Callithrix jacchus*) are the most common non-human primates (NHPs) used in these investigations. Since great cost is associated with developing and maintaining a colony of NHPs, early research and toxicological studies are usually carried out in rodent models. In studies using behavioral measures such as drug addiction, NHPs are the first choice test animal. In addition, NHPs and individual humans exhibit differential sensitivities to many drugs and toxins and can be

classified as "extensive metabolizers" and "poor metabolizers" of these agents.

In additional embodiments, the nucleic acid molecules which encode the mammalian protein may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleic acid molecules that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

### Examples

It is understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary. It is also understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. The examples below are provided to best describe the subject invention and its representative constituents.

#### I cDNA Library Construction

The RALINOT01 cDNA library was constructed from liver tissue removed from a pool of fifty 10- to 11-week-old Sprague-Dawley female rats (Pharmakon, Waverly PA). The animals were housed in standard laboratory caging and fed PMI-certified Rodent Diet #5002. The animals appeared to be in good health at the time tissue was harvested. The animals were anesthetized by CO<sub>2</sub> inhalation, and then cardiocentesis was performed.

Frozen tissue was homogenized and lysed in TRIZOL reagent (1 g tissue/10 ml TRIZOL; Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate, using a POLYTRON homogenizer (PT-3000; Brinkmann Instruments, Westbury NY). After a brief incubation on ice, chloroform (1:5 v/v) was mixed with the reagent, and then centrifuged at 1,000 rpm. The upper aqueous layer was removed to a fresh tube, and the RNA precipitated with isopropanol, resuspended in DEPC-treated water, and treated with DNase I for 25 min at 37°C. The RNA was re-extracted once with phenol-chloroform, pH 4.7, and precipitated using 0.3 M sodium acetate and 2.5 volumes ethanol. The mRNA was then isolated using an OLIGOTEX kit (QIAGEN, Chatsworth CA) and used to construct the cDNA library.

The mRNA was handled according to the recommended protocols in the SUPERScript plasmid system (Life Technologies). The cDNAs were fractionated on a SEPHAROSE CL-4B column (Amersham Pharmacia Biotech), and those cDNAs exceeding 400 bp were ligated into the pINCY1 plasmid vector (Incyte Pharmaceuticals). The plasmid pINCY1 was subsequently transformed into DH5α or DH10B competent cells (Life Technologies).

The RAKINOT01 library was constructed using mRNA isolated from kidney tissue removed from a pool of fifty, 7- to 8-week-old male Sprague-Dawley rats, as described above.

The RAKINOT02 library was constructed using mRNA isolated from kidney tissue removed from a pool of fifty, 10- to 11-week-old female Sprague-Dawley rats, as described above.

## II cDNA Library Normalization

In some cases, cDNA libraries were normalized in a single round according to the  
5 procedure of Soares et al. (1994, Proc. Natl. Acad. Sci. 91:9228-9232) with the following  
modifications. The primer to template ratio in the primer extension reaction was increased from  
2:1 to 10:1. Reduction of each dNTP concentration in the reaction to 150  $\mu$ M allowed the  
generation of longer (400-1000 nucleotide (nt)) primer extension products. The reannealing  
hybridization was extended from 13 to 19 hours. The single stranded DNA circles of the  
10 normalized library were purified by hydroxyapatite chromatography, converted to partially double-  
stranded by random priming, and electroporated into DH10B competent bacteria (Life  
Technologies).

The Soares normalization procedure is designed to reduce the initial variation in individual  
cDNA frequencies and to achieve abundances within one order of magnitude while maintaining  
15 the overall sequence complexity of the library. In the normalization process, the prevalence of  
high-abundance cDNA clones decreases significantly, clones with mid-level abundance are  
relatively unaffected, and clones for rare transcripts are increased in abundance. In the modified  
Soares normalization procedure, significantly longer hybridization times are used to increase gene  
discovery rates by biasing the normalized libraries toward low-abundance cDNAs that are well  
20 represented in a standard transcript image.

The RALINON03, RALINON04, and RALINON07 normalized rat liver cDNA libraries  
were constructed with  $2.0 \times 10^6$ ,  $4.6 \times 10^5$ , and  $2.0 \times 10^6$  independent clones from the RALINOT01  
cDNA library, respectively. The RALINOT01 cDNA library was normalized in one round using  
conditions adapted from Soares (supra) except that a significantly longer (48-hour) reannealing  
25 hybridization was used.

## III cDNA Library Prehybridization

The RALINOH01 cDNA library was constructed with clones from the RALINOT01  
cDNA library. After preparation of the RALINOT01 cDNA library, 9,984 clones were spotted  
onto a nylon filter, lysed, and the plasmid DNA was bound to the filter. The filter was incubated  
30 with pre-warmed hybridization buffer and then hybridized at 42°C for 14-16 hours in 0.75 M  
NaCl, 0.1 M  $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$ , 0.15 M tris-HCl (pH 7.5), 5x Denhardt's Solution, 2% SDS, 100  
 $\mu$ g/ml sheared salmon sperm DNA, 50% formamide, and [ $^{32}$ P]-labeled oligonucleotide molecules  
made from reverse transcribed rat liver mRNA from an untreated animal. The filter was rinsed  
with 2 x SSC (saline sodium citrate) at ambient temperature for 5 minutes followed by washing for  
35 30 minutes at 68°C with pre-warmed washing solution (2 x SSC, 1% SDS). The wash was

repeated with fresh washing solution for an additional 30 minutes at 68°C. Filters were then washed twice with pre-warmed washing solution (0.6 x SSC, 1% SDS) for 30 minutes at 68°C. Some 4,224 clones had very low hybridization signals and about 20% of the clones had no signals and two groups were isolated and sequenced.

5     **IV     Isolation and Sequencing of cDNA Clones**

DNA was isolated using the following protocol. Single bacterial colonies were transferred into individual wells of 384-well plates (Genetix Ltd, Christchurch, United Kingdom) using sterile toothpicks. The wells contained 1 ml of sterile Terrific Broth (Life Technologies) with 25 mg/l carbenicillin and 0.4% glycerol (v/v). The plates were covered and placed in an incubator  
10 (Thermodyne, Newtown Square PA) at 37°C for 8-10 hours. Plasmid DNA was released from the cells and amplified using direct link PCR (Rao, V.B. (1994) Anal. Biochem. 216:1-14) as follows. The direct link PCR solution included 30 ml of NUCLEIX PLUS PCR nucleotide mix (Amersham Pharmacia Biotech, Piscataway NJ) and 300 µl of Taq DNA polymerase (Amersham Pharmacia Biotech). Five microlitres of the PCR solution were added to each of the 384 wells using the  
15 MICROLAB 2200 system (Hamilton, Reno NV); plates were centrifuged at 1000 rpm for 20 seconds and refrigerated until use. A 384 pin tool (V&P Scientific Inc, San Diego CA) was used to transfer bacterial cells from the incubation plate into the plate containing the PCR solution where 0.1% Tween 20 caused the cells to undergo lysis and release the plasmid DNA. After lysis, the plates were centrifuged up to 500 rpm, covered with a cycle sealer, and cycled using a 384-  
20 well DNA ENGINE thermal cycler (MJ Research, Watertown MA) using the program dPCR30 with the following parameters: Step 1) 95°C, 1 minute; Step 2) 94°C, 30 seconds; Step 3) 55°C, 30 seconds; Step 4) 72°C, 2 minutes; Step 5) steps 2, 3, and 4 repeated 29 times; Step 6) 72°C, 10 minutes; and Step 7) storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 µl PICO  
25 GREEN quantitation reagent (0.25% (v/v), Molecular Probes, Eugene OR) dissolved in 1x TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the quantitation reagent. The plate was scanned in a Fluoroscan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantitate the concentration of DNA. Typical concentrations of each DNA sample were in the  
30 range of 100 to 500 ng/ml.

The cDNAs were prepared for sequencing using either a HYDRA microdispenser (Robbins Scientific, Sunnyvale CA) or MICROLAB 2200 system (Hamilton) in combination with the DNA ENGINE thermal cyclers (MJ Research). The cDNAs were sequenced using the method of Sanger, F. and A.R. Coulson (J. Mol. Biol. (1975) 94:441-448) and the ABI 377 sequencing  
35 systems (PE Biosystems). Most of the isolates were sequenced according to standard ABI

protocols using ABI kits (PE Biosystems). The solution volumes were used at 0.25x - 1.0x concentrations. Typically, 500 to 700 base pairs were sequenced in 3.5 to 4 hours. In the alternative, cDNAs may have been sequenced using solutions and dyes from Amersham Pharmacia Biotech.

5     **V     Rat Liver and Kidney Gene Selection**

As a first step, originator molecules from high throughput sequencing experiments were derived from clone inserts from RALINOT01, RAKINOT01, RAKINOT02, RALINOH01, RALINON03, RALINON04 and RALINON07. cDNA library clones were obtained. There were 18,140 rat liver molecules and 5,779 rat kidney molecules.

10     Additionally, 1,500 rat molecules derived from clone inserts of any of 113 rat cDNA libraries were selected based on their homology to genes coding for polypeptides implicated in toxicological responses including peroxisome-associated genes, lysosome-associated genes, apoptosis-associated genes, P450 cytochromes, detoxification genes such as sulfotransferases, glutathione S-transferase, and cysteine proteases, and the like.

15     Then, all the remaining molecules derived from all of the rat cDNA library clones were clustered based on the originator molecules described above. The clustering process involved identifying overlapping molecules that have a match quality indicated by a product score of 50 using BLAST. 6581 master clusters were identified.

20     After forming the clone clusters, a consensus sequence was generated based on the assembly of the clone molecules using Phrap (Phil Green, University of Washington). The assembled molecules were then annotated by first screening the assembled molecules against GenBank using BLASTn and then by screening the assembled molecules against GenPept using FASTX. About two thirds of the assembled molecules were annotated, about one third of the  
25     assembled molecules were not annotated.

**VI     Substrate and Array Element/Probe Preparation**

Clones nominated in the process described in Example V were used to generate array elements. Each array element was amplified from bacterial cells. PCR amplification used primers complementary to the vector sequences flanking the cDNA insert. Array elements were amplified  
30     in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 µg. Amplified array elements were then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements were immobilized on polymer-coated glass slides. Glass microscope slides (Corning, Corning NY) cleaned by ultrasound in 0.1% SDS and acetone, with  
35     extensive distilled water washes between and after treatments. Glass slides were etched in 4%

hydrofluoric acid (VWR, West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma-Aldrich, St. Louis MO) in 95% ethanol. Coated slides were cured in a 110°C oven.

Array elements were applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522 and incorporated herein by reference. In brief, 1 µl of the array element DNA, at an average concentration of 0.5 µg/ml in 3 x SSC, was loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposited about 5 nl of the array element sample per slide. A total of 7404 array elements representing rat liver and kidney genes and a variety of control elements, including 14 synthetic control molecules, human genomic DNA, and yeast genomic DNA, were arrayed in four identical quadrants within a 1.8 cm<sup>2</sup> area of the glass substrate.

Microarrays were UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays were washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites were blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

## VII Target Preparation

Male Sprague-Dawley rats (6-8 wk old) were dosed intraperitoneally with clofibrate (CLO; Acros, Geel, Belgium) at 250 mg/kg body weight (bw), acetaminophen (APAP; Acros) at 1000 mg/kg bw, benzo(a)pyrene (B(a)P; Acros) at 10 mg/kg bw, or dimethylsulfoxide vehicle (DMSO; Acros) at less than 2 ml/kg bw and the animals were later euthanized by CO<sub>2</sub> inhalation. Animals were monitored daily for physical condition and body weight. Three animals per group were sacrificed approximately 12 hours, 1 day (d), 3d, 7d, 14d, and 28d following the single dose. Prior to sacrifice a blood sample from each animal was drawn and assayed for serum alanine transferase (ALT) and aspartate aminotransferase (AST) levels using a diagnostic kit (Sigma-Aldrich). Observed gross pathology and liver weights were recorded at time of necropsy. Liver, kidney, brain, spleen and pancreas from each rat were harvested, flash frozen in liquid nitrogen, and stored at -80°C.

For each probe preparation, frozen liver was homogenized and lysed in TRIZOL reagent (Life Technologies, Gaithersburg MD) following the modifications for liver RNA isolation. Messenger RNA was isolated using an OLIGOTEX kit (QIAGEN) and labeled with either Cy3- or Cy5-labeled primers (Operon Technologies, Alameda CA) using the GEMBRIGHT labeling kit (Incyte Pharmaceuticals). Messenger RNA isolated from tissues of rats treated with clofibrate, acetaminophen, or benzo(a)pyrene was labeled with Cy5 and mRNA isolated from tissues of rats treated with DMSO was labeled with Cy3. Quantitative and differential expression pattern control

cDNAs were added to each labeling reaction. Labeled cDNA was treated with 0.5 M sodium bicarbonate (pH 9.2) for 20 min at 85°C to degrade the RNA and purified using two successive CHROMA SPIN 30 gel filtration spin columns (Clontech, Palo Alto CA). Cy3-labeled control sample and Cy5-labeled experimental sample were combined and precipitated in glycogen, sodium acetate, and ethanol.

Probes are also prepared from tissue needle biopsy samples. Samples are used to identify changes within the tissue following exposure to, for example, a toxic compound, a potential toxic compound, a compound with unknown metabolic responses, or a pharmacological compound.

### VIII Hybridization

Hybridizations were carried out using the methods described by Shalon (supra).

### IX Detection

The microscope used to detect the reporter-labeled hybridization complexes was equipped with an Innova 70 mixed gas 10 W laser (Coherent Lasers, Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3, and 632 nm for excitation of Cy5. The excitation laser light was focused on the array using a 20x microscope objective (Nikon, Melville NY). The slide containing the array was placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example was scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excited the two fluorophores sequentially. Emitted light was split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics, San Jose CA) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes were used to filter the signals. The emission maxima of the fluorophores used were 565 nm for Cy3 and 650 nm for Cy5. Each array was typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus was capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans was typically calibrated using the signal intensity generated by a cDNA control species added to the probe mix at a known concentration. A specific location on the array contained a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two probes from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration was done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube was digitized using a 12-bit RTI-835H analog-to-

digital (A/D) conversion board (Analog Devices, Norwood MA) installed in an IBM-compatible PC computer. The digitized data were displayed as an image where the signal intensity was mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data was also analyzed quantitatively. Where two different

5 fluorophores were excited and measured simultaneously, the data were first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid was superimposed over the fluorescence signal image such that the signal from each spot was centered in each element of the grid. The fluorescence signal within each element

10 was then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis was the GEMTOOLS gene expression analysis program (Incyte Pharmaceuticals). In one analysis, where two different samples were prepared from identically treated cell cultures, expression patterns of those cDNAs which changed between 1.6- and 1.7-fold were within the 95% confidence limits of a Poisson normal distribution profile

15 (T. Theriault, pers. communication).

## X Results

The expression patterns of eight cytochrome P450 isozymes known to be induced in a toxicological response were monitored during the 28 day time course. The results using clofibrate, acetaminophen, and benzo(a)pyrene are shown in Table 1, Table 2, and Table 3, respectively.

20 Each of the known genes was upregulated greater than 2 fold at least once during the time course.

TABLE 1 Gene expression patterns (x fold increase) of known genes in clofibrate-treated rat liver

Gene	12 hours	1 day	3 days	7 days	28 days
P450 LA- omega	15	26	2.0	2.1	3.0
P450 4A	6.5	16.5	2.1	3.0	3.5
P450 3A	0.14	1.6	0.63	0.50	0.45

TABLE 2 Gene expression patterns (x fold increase) of known genes in acetaminophen-treated rat liver

Gene	12 hours	24 hours	3 days	7 days	14 days	28 days
P450A	1	4.4	2.2	2.0	4.6	4.8
P450F	0.50	0.23	2.0	1.8	2.2	2.2
P450 14DM	0.45	0.32	2.2	1.6	1.8	0.56



TABLE 3 Gene expression patterns (x fold increase) of known genes in benzo(a)pyrene-treated rat liver

Gene	12 hours	1 day	3 days	7 days	14 days	28 days
P450 LA-omega	1.2	2.3	2.4	1.4	6.8	1.2
P450 MCA-inducible	8.2	11.8	4.4	2.2	2.4	1.2
P450 ISF/B-NF	9.6	7.4	6.2	2.4	2.4	1.2

We have discovered novel nucleotide molecules that are up-regulated or down-regulated at least 2-fold at least once during the time course. These molecules are SEQ ID NOs:1-117 provided in the Sequence Listing. These polynucleotide molecules can be used for screening test compounds or molecules for a toxicologic effect.

Table 4 shows the gene expression pattern of selected molecules that were upregulated at least 2-fold at least once during the time course following treatment with clofibrate (CLO) and Table 5 shows the gene expression pattern of selected molecules that were downregulated at least 2-fold at least once during the time course following treatment with clofibrate.

TABLE 4 Gene expression patterns (x fold increase) of CLO-upregulated nucleic acid molecules

SEQ ID NO:	12 hours	1 day	3 days	7 days	28 days
35	11.6	14.4	2.4	3.0	3.2
36	11.6	18.7	3.0	3.3	3.8
31	1.2	2.8	1.0	2.3	4.8
57	0.9	1.9	0.9	1.5	4.5
67	4.3	1.1	1.6	1.7	5.7
81	5.1	1.2	1.7	1.8	6.0
94	4.8	1.4	2.0	1.5	2.4
33	5.1	1.3	1.9	1.8	5.5



TABLE 5 Gene expression patterns (x fold increase) of CLO-downregulated nucleic acid molecules

SEQ ID NO:	12 hours	1 day	3 days	7 days	28 days
102	0.15	1.4	1.0	0.77	0.67
103	0.13	1.2	0.83	0.63	0.56
52	0.13	0.56	0.37	1.0	1.2
43	0.13	1.1	0.91	0.71	0.56
53	0.11	0.67	0.36	1.0	1.2
54	0.14	0.63	0.59	1.1	0.29
55	0.16	0.67	0.71	1.2	0.32
63	0.33	0.14	1.1	0.83	1.2
105	0.14	1.2	1.0	0.77	0.71
68	0.16	0.67	0.53	1.1	1.4
71	0.43	0.18	0.40	0.34	0.23
74	0.06	0.71	0.42	1.1	1.2
115	0.22	1.5	0.77	1.7	1.3
85	0.19	0.45	1.0	1.3	1.8
90	0.12	0.48	1.2	1.0	1.2
95	0.14	0.91	0.56	1.5	1.4

Table 6 shows the gene expression pattern of selected molecules that were upregulated at least 2-fold at least once during the time course following treatment with acetaminophen (APAP) and Table 7 shows the gene expression pattern of selected molecules that were downregulated at least 2-fold at least once during the time course following treatment with acetaminophen.

TABLE 6 Gene expression patterns (x fold increase) of APAP-upregulated nucleic acid molecules

SEQ ID NO:	12 hours	24 hours	3 days	7 days	14 days	28 days
35	3.1	6.6	2.9	3.3	4.9	7.5
36	4.7	10.1	4.0	4.2	6.9	9.8
78	0.9	4.4	1.2	1.5	1.1	1.4
81	2.9	5.1	1.4	1.8	2.3	2.4
82	1.2	4.2	1.3	1.0	1.7	1.4
39	2.4	9.0	2.6	1.7	2.2	2.4
94	1.2	4.9	1.2	1.1	2.0	2.0
33	4.3	5.9	1.5	1.7	2.9	3.2
98	1.3	6.1	1.5	1.9	1.8	2.1

TABLE 7 Gene expression patterns (x fold increase) of APAP-downregulated nucleic acid molecules

SEQ ID NO:	12 hours	1 day	3 days	7 days	14 days	28 days
49	0.59	0.15	1.2	1.0	0.83	1.1
50	0.83	0.37	0.43	0.37	0.22	0.2
52	0.63	0.08	1.0	0.71	0.83	0.45
53	0.25	0.07	1.1	0.71	0.83	0.42
54	0.43	0.19	0.04	0.71	0.29	0.36
55	0.35	0.22	0.07	0.77	0.31	0.42
56	0.38	0.21	0.5	0.32	1.1	1.1
59	0.18	0.77	2.5	1.4	1.2	1.6
61	0.15	0.53	0.91	0.71	0.71	1.8
63	0.13	0.05	0.23	0.77	0.43	0.77
74	0.19	0.09	1.1	1.0	1.4	0.56
87	0.42	0.10	0.53	0.63	0.63	0.67
90	0.16	0.29	1.2	0.77	0.83	1.1
95	0.22	0.20	2.7	1.7	1.6	1.0

Table 8 shows the gene expression pattern of selected molecules that were upregulated at least 2-fold at least once during the time course following treatment with benzo(a)pyrene (B(a)P) and Table 9 shows the gene expression pattern of selected molecules that were downregulated at least 2-fold at least once during the time course following treatment with benzo(a)pyrene.

5

TABLE 8 Gene expression patterns of B(a)P-upregulated nucleic acid molecules

SEQ ID NO:	12 hours	1 day	3 days	7 days	14 days	28 days
3	3.4	1.9	0.7	0.5	1.99	0.77
9	1.6	3.2	1.2	1.1	3	1.5
10	2.8	5.9	3.2	2.1	2.9	1.8
13	2.9	6.1	3.1	2.3	3.3	1.9
19	2.7	3.5	3	1.9	1.7	1.5
26	1.1	4.7	1.5	1.3	5	2
31	2.3	3.8	1.6	2	1.7	2.1
33	2.1	4.1	3.2	2	1.7	1.6
35	1.2	3	5.1	1.4	5	1.3
37	3.4	0.5	0.6	0.7	0.9	0.5
39	1.5	3.5	1.8	1.5	3.5	2.1
42	9.1	9.1	5.2	2.4	2.1	1

TABLE 9 Gene expression patterns of B(a)P-downregulated nucleic acid molecules

SEQ ID NO:	12 hours	1 day	3 days	7 days	14 days	28 days
11	0.3	0.5	0.4	0.3	0.53	0.53
25	0.3	0.9	0.5	0.7	0.42	2.1
27	1	0.1	1	1.1	0.09	0.53
28	0.3	0.3	1.2	1.2	0.77	1.1
45	1.2	0.2	0.4	0.6	0.77	0.37

30

## CLAIMS

What is claimed is:

1. A method for detecting or diagnosing the effect of a test compound or molecule associated with increased or decreased levels of a nucleic acid molecule in a mammalian subject comprising:
  - 5 a) treating a mammalian subject with a toxic compound or molecule;
  - b) obtaining a sample containing nucleic acids from the mammalian subject treated with the toxic compound or molecule;
  - c) contacting the sample with a microarray comprising a plurality of nucleic acid molecules comprising SEQ ID NOs: 1-117, or a fragment thereof, under conditions for the
  - 10 formation of one or more hybridization complexes;
  - d) detecting the hybridization complexes, wherein the presence, absence or change in amount of the hybridization complex, as compared with the hybridization complexes formed from nucleic acid molecules from an untreated mammalian subject, is indicative of a metabolic response to the toxic compound or molecule;
  - 15 e) measuring the level of nucleic acid molecules in a sample from a mammalian subject treated with a test compound or molecule using the method of steps (c) and (d); and
  - f) comparing the level detected in step (e) to a level of nucleic acid molecules present in normal or untreated biological sample in which an increase or decrease in the level of nucleic acid molecule as compared to normal levels indicates a toxicological response.
- 20 2. The method of claim 1 wherein the toxic compound or molecule is selected from hypolipidemic drugs, n-alkylcarboxylic acids, n-alkylcarboxylic acid precursors, azole antifungal compounds, leukotriene D4 antagonists, herbicides, pesticides, phthalate esters, phenyl acetate, dehydroepiandrosterone sulfate, oleic acid, methanol and their corresponding metabolites, acetaminophen and its corresponding metabolites, benzo(a)pyrene, 3-methylcholanthrene,
- 25 benz(a)anthracene, 7,12-dimethylbenz(a)anthracene, and their corresponding metabolites.
3. The method of claim 1 wherein the sample is a tissue selected from the group consisting of liver, kidney, brain, spleen, pancreas, and lung.
4. The method of claim 1 wherein the test compound which elicits the metabolic response is a compound with previously unknown metabolic response.
- 30 5. The method of claim 1 wherein the test compound or molecule which elicits the metabolic response induces at least a 2-fold change in the amount of the hybridization complexes formed with at least one of the nucleic acid molecules of the sample.
6. An isolated and purified nucleic acid molecule selected from SEQ ID NOs: 1-117, or a fragment thereof.
- 35 7. A method of using the nucleic acid molecule of claim 6 to screen a library of molecules or

compounds to identify at least one molecule or compound which specifically binds the nucleic acid molecule, the method comprising:

- a) combining the nucleic acid molecule of claim 6 with a library of molecules or compounds under conditions to allow specific binding; and
  - 5           b) detecting specific binding, thereby identifying a molecule or compound which specifically binds the nucleic acid molecule.
8.       The method of claim 7 wherein the library is selected from DNA molecules, RNA molecules, peptide nucleic acids, artificial chromosome constructions, peptides, and proteins.

## SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.  
CUNNINGHAM, Mary Jane  
ZWEIGER, Gary B.  
PANZER, Scott R.  
SEILHAMER, Jeffrey J.

<120> TOXICOLOGICAL RESPONSE MARKERS

<130> PA-0010 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/141,825; 09/172,711; 09/172,108

<151> 1998-08-28; 1998-10-13; 1998-10-13

<160> 117

<170> PERL Program

<210> 1

<211> 259

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700416862F7

<400> 1

gtggcggcga tttctgcgtc gagcatttgg agtttcttcg ctgctgaacg ggtagactaa 60  
acggcggtcg acatggtgga ggaggtacag aagcattctg tgcacacact agtgttcagg 120  
tcattgaaga ggacccatga catgtttgtg gctgataatg gaaaacctgt gcctttggat 180  
gaagagagtc acaagcggaa aatggcaatc aagcttcgta atgagtatgg ccctgtgctg 240  
catatgccta cttcaaaag 259

<210> 2

<211> 295

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 212, 227, 229, 232, 240, 243, 245, 250, 257, 267, 269, 273, 277

<222> 288, 290

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700502957F6

<400> 2

gcctcttcca ccatccggcc tagtcaactgc aggggccatg cctacctatt ccaactcaact 60

tgttacctct gcggctccag gcagggctta gtccaacctg cccagacacg gttcaccttt 120  
 ttatgcccac gctttcgggg tgctgaggta ggggctgcct tcctgcaccc ccaaggagca 180  
 gacactcaag aatggagtca gctaggaacc cngggagctg cctcatnang cnettgatan 240  
 cangnacacn tttgcanctg cagacctntt tcngaanaac nttgccangn tcaac 295

<210> 3  
 <211> 273  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 56, 220, 235, 237, 239, 249, 256  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700504502F6

<400> 3  
 tgacctgct tgctgcaggt gactgggtcaa gtgcgagcta gcttagttag tgtgngtat 60  
 aaggcgccat cattcctcca gtaagcctcc atcccaaagc aactgaggct gtggcagtga 120  
 tgccagcaac ctgtgtcacc caaaattatc cagccctcca cgggcaactgc ctaggacctg 180  
 gggagggaag ggactttgca tcacatagcc tcagggttcgn gtttggtctt ggtangngnt 240  
 gcctgaaant ggtggnttcc agctggtgta cgg 273

<210> 4  
 <211> 264  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700505170F6

<400> 4  
 ggctggctctg cgatggcccc ctacctcggc tcgctggaac catgtgtggg tccggcactt 60  
 gagactggaa tcctgaaagg ggtgaacctt cagcggaaac ttgcggcaaa ttttactccg 120  
 tccggacagc cacggcgagg ggaggcagtg aatgctttgt gctggggcac aggcggcgag 180  
 acccagattt tgggtgggatg tgccgacagg accgtgaggc actttaatgc ggaggagggt 240  
 acatccaga ccagagatac tgcc 264

<210> 5  
 <211> 268  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 231  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature



<223> Incyte template ID No: 700511170F6

<400> 5

```
ggcaacaaga cgctgtgatt ggaagcaatg acgaagtcct cactctcgg ggagtgggta 60
tgtgtgcta catcatgtga tgggcagcct ggaggggatg caggcgccct ggagctatgt 120
ccaggtggc atgggtgccc tctcagatgc cattgcaagc tcggctactg cacatggagc 180
aagtatcttc acagagaaga ctgtgggctaa ggtgcaagtg aacagcgaag ncgtgtccaa 240
ggggtcgtgc tcagggccgg cgaggagt 268
```

<210> 6

<211> 284

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 28, 37, 44, 67, 71, 88, 102, 172, 238

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700607827F7

<400> 6

```
gtgtacctac ctgctgagga aggaaggngt ggatggnaacc cagnaacctg atgtccagca 60
caagggnaac nggcgtggct ttctaantna caagcctggg tnactacagc ttgcagctac 120
ctaaccatg caggaggcga accctctgag ccagttgct attgtgacca tnaagatgtc 180
ttgccacaca gcttccaccc agtctgggtt taatgggaag ttacctaacg attaccnca 240
gaagacacat gagacgcttg cttcaaagct ctcggatgca gccca 284
```

<210> 7

<211> 243

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700500814F6

<400> 7

```
actttttcta atgtcttatg gccacttctt atgagaatgg ggagctgtc tgctgaggg 60
tcgtgagagg aagcgccaga gcaggcccat catccaacc cttggccttg gcccttccc 120
ctagctctgc agcatttctt cagatcctct ttctgagag tcaaggagac taaacaccaa 180
taaaccagac acaaccttcg tggcccaaaa ggagaaaccg attagagggg tctctgctag 240
atg 243
```

<210> 8

<211> 259

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 235

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700526819F6

<400> 8

```
ggcgccggcg gaaggggcct gtcaccgtcc gctgcgacgt cgcggctgga gttgaacctg 60
gtgccggctg ctttgcgctg tgagtcgatg gcggccgaaa aacgagaacc ggacgagtgg 120
cgtctggaga agtatgtggg tccctggaag acatgctgca gccctgaaag tccaagcaag 180
taaaccgcct cggaagtgat cagtgagtac tcccgcgaaag tgactttctg aaggngatgct 240
gaggctgaga agtgactct 259
```

<210> 9

<211> 255

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 2, 4, 53, 55, 86, 88, 92, 106, 133, 211, 214

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700528176F6

<400> 9

```
cnanggccct caggggaatca agaggagcca gcctgatccc tggcctctgg agncntaaaa 60
caagtgtgtt tttgcaggta gcctangntg gntgtcgatg gagctncagc ctgcatggca 120
ttaggcagga agncaactctg gatgattgtg cacatgagaa cctagtcagg gagggagggt 180
ttaaggagag gcttagaata caagtgagaa ncancgagaa agggaccaag tcctcagaat 240
agaagctatc tgcct 255
```

<210> 10

<211> 269

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 30-31, 140

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700528082F6

<400> 10

```
ataaaagtga aaactgggca agggcaaggng ngctgggcgt gaaccgctta ctagataatg 60
gtctctaaaa attggctctg aaaaccctgt ttgtgtattc gttttatgag tgcttaaaaa 120
tggtgtgacc agggcatggn cactgtcatt ggaacagcaa catgcttgct ggcacattgg 180
aatgggggaaa tgtgaagaaa gctggcatca ggctgcggc acccatttct ttgatgaaag 240
tggtgtgtca aacccccact aatcatttt 259
```

<210> 11  
<211> 254  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 36, 47, 67, 80, 82-83, 92, 111  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700535328F6

<400> 11  
caaggcggttc tgctgcgaga acgacattga catcgngcgc gtggtanacg tgcggaggct 60  
ggcggcnatc gtggcgccn annacgagag gngcgcccg tgagacttgc nttgcatcct 120  
catttcgaac cctaataag acacatggaa ggacctgcc ttggagaagc tcagtttggt 180  
ctgcgaggag agccgcagct tcaacgactg ggtcccagca tcaccttcc gactgacagc 240  
ctgcaggac cttg 254

<210> 12  
<211> 244  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 50, 102, 169, 171, 204, 210, 234, 242  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700368973F6

<400> 12  
cacaatccca aactggaaaa acttaaaaag gaatcctgct gtgaaaggtn tatattactc 60  
tagatttttc ttactgtaaa tattgtaaga ttgtaatact gncaatattt tattaaccaa 120  
caaagttaa tctatgtgaa atcagactta tttaaagggc tgctattang ngtgtggccc 180  
tttgctgaca gattaagtat attntgagtn agataactta ttaaggatgg aacnttaaag 240  
gntc 244

<210> 13  
<211> 237  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 11, 20, 132, 136, 149, 151, 158, 171, 173, 184, 187, 190, 193  
<222> 204-205, 208, 212-213, 215, 221, 227, 230-231, 236  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature

<223> Incyte template ID No: 700368974F6

<400> 13

```
gtccttagct ngtgggcggn ggggtgcagt ctgctactat ctgaacgaat ttaatgtggg 60
agcatgcctt atacaacaca ggaaattaat gtgtgatcta atgcgtgac tatgacttat 120
tacaatacag anttangtgt gaacctgcnt ncaaaacngg tcagaatttt ngnaatggcc 180
ggantgnacn ggntgnttat taanntgnaa gnnngntggga naggccnggn ntgcgnt 237
```

<210> 14

<211> 235

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 38, 49, 53, 55, 107, 125, 132, 136, 138, 150-151, 163, 165-166

<222> 179-180, 184, 186, 189, 192, 199, 219-221, 223

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700369461F6

<400> 14

```
ctttaaccgg tgggctgctg taagaatcgg tggcaggnet ctctcggg ggngntaatt 60
gctctggaac gctactagga cccgaataact aaggccacat ctctacgic taagagggga 120
aatangatag cnttgnntcc acatgtggcn nagtgggggt gcngnntatn gcttaacann 180
tactnantnc antgattant gtggtggtaa gatggcttnn ntnaaaactg ccgcc 235
```

<210> 15

<211> 205

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 17, 19, 21, 56, 69, 75, 81, 103, 109, 111, 122, 127, 130, 137

<222> 148, 151, 153, 166, 174, 177, 190, 193, 195

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700782579F6

<400> 15

```
gggtcattta caacctntna nacaagggga cgccccaga catgccagtg ttcacngaac 60
aagatgcang tccancagga ncagatagac tcagtccatg gantggctna ncaaccagcc 120
cngggcngcn caactgntgg acangtangg nantttctgc gacaantggt gagnttnttg 180
cgaagctccn tgnengaaag aatgg 205
```

<210> 16

<211> 236

<212> DNA

<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 11-12, 33-34, 46, 76, 85-86, 103, 123, 127, 144, 152, 157, 162  
<222> 169, 174, 183-184, 186, 191-194, 217, 224, 230  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700782665F6

<400> 16  
ggctccatg nngaccgggc gccttggggt tgnngagacg ctgcangccc ttaacgccc 60  
cttgtagggc ccttgnaacc ccggnntggg ttaaggaaaa cgnatgccc acaccttcgt 120  
gtncganact ttttggcacc gcgnaaactc gnntacngtg gnttttnang tagnggggtat 180  
gtnnncgag nnnnttttagg ccggcttggt ctgcggnatt gggntgaaan tgtctg 236

<210> 17  
<211> 267  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700480927F6

<400> 17  
ctggaccaac atcacaagaa atgaataaag cagatttctc tgttgagttc tgcagtaaac 60  
cacctaaaag ccaatgtcaa gtcagccgca gacttactta gcctgectag cactgtagag 120  
ggacttcaga agagtgtcgc ttccattggc aatacgttga acagtgtcag ccttgctgta 180  
gaggcaatac agaagaccgt ggatgaacac aaggcacctt ggagttactg cagggcagtg 240  
tggagaccaa tgggaagcaac caaatca 267

<210> 18  
<211> 271  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 2, 92, 162, 181, 247-248, 253, 256, 262, 269  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700481732F6

<400> 18  
gncttattta tgtatgaaaa tgcagaaatc tgtacattcc tcaagccagt cctgtcgagc 60  
caggtctgtc ccatacctgt acctcaaccc antcccacct ggctgaaca cccatgaga 120  
cagagctggg ctctgggctg gggccccag gcctgggctg gncaggcaga cctaccccc 180  
nagtccactg gctccagtct ccgaggctct cctgggctac aaagggggac cacacacacc 240  
cagaatnntt tantgnattg gngggccng g 271

<210> 19

<211> 283  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 37, 131, 199, 205-206, 208, 232-233, 239, 244-246, 251, 253-254  
 <222> 261-262, 266, 269, 274-275, 280  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700483143F6

<400> 19  
 ctaaaattaa gatagaagtg aatgagacag atatctngta agacactgta ttttcttgtg 60  
 tgatcagatc tagtgtgtg ggatgataga agttgaactt gctttattgc tatgggttaa 120  
 aatattttgt ntcattaaaa tggcctattg aaatgctttt ctgttcctat aataaaataa 180  
 cctgatgaaa aagtaaaana aaaannanaa aaaaaaagg gcggtccccg cnnagggcnt 240  
 tttnnncccg ngnttantt nngcnggnc cctnnggggn cca 283

<210> 20  
 <211> 256  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700484538F6

<400> 20  
 ctggcctcag cttcctaagt tctgggagtc ggacagggtg ttgccacaca catggcccac 60  
 cggggacctga gaacctacag tgaaccgtca ccagggtct gtggatgttc tgcattctga 120  
 ggtagacagc ctctaatac ctgttaggga cctaggacca gagctggggt gccaggcat 180  
 gtcccaacat gtgcctcgg ccacagggat atcggttgaa gtgcatttgg aagtgtgctg 240  
 ggaagccagc cagctt 256

<210> 21  
 <211> 272  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 64, 68, 124, 140, 145, 213, 260, 267, 269  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700480077F6

<400> 21  
 ctgacctgac ccatgattta aggaccgtag tttagcacgg accactgcaa aggggggcta 60  
 aggnctgntg ggctaaaggt ctctttgagc ccagtggtta tagtcacacc ttctttgctc 120  
 tggncaggga ggcctacttn ttctntactc gtggaatcct ggaatcttaa agataaaaga 180

acctagaaag aaaatcaaac ccactttcct tgnnggcaga tggtaatatg ggactgagac 240  
 agcaaacctg gggcttggan aggaccnanc tc 272

<210> 22  
 <211> 270  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 40, 44, 46, 114  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700060207F7

<400> 22  
 tacccttgca tgggataccg tttctcgacc ccagtgcacn tggntnctgt catcccatga 60  
 aattgcagca agggcagctct cttttgtggg aacagattaa ctccctacaca tgangtagat 120  
 tcaacacctg ccaggaaagc agaagcatta ctttaagtgtc ctgtgaaggc aaacatcaag 180  
 tcaattcagc ttatcttgaa gagtggcaaa ccatgaactc caaatgtcat tgtgtgaaac 240  
 tgaacgatgg tcatttcatt ccggtgctgg 270

<210> 23  
 <211> 250  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 45, 49-50  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700133105F6

<400> 23  
 ccctcctgta tctgaaccca gcttctcagc tctgagatga gtgcnggann ggcttcccaa 60  
 cctatgotca ataccacagg cagcctgcag gaggagaaaa tgggtaaaat gttccatggg 120  
 aaatgtctca gaatcgtctc ccccgaaatct cctgctaagc tttactgctg ctatggagtg 180  
 atcatgggtcc tcagtgtagc tgtagttgct ctttctgttg ctttgtcagt aaaaatgaca 240  
 ccacagatct 250

<210> 24  
 <211> 226  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700138117F6

&lt;400&gt; 24

gaggattcac tcacatttgc ttcccgtggt ccatgagtga gctgcccttt ctgagtcag 60  
agggagccag agggcctcac aacaacagag ggtctcagag ctccctggag gaaggctcag 120  
ttacaggctc agaggctcgg cacagcttag gtgtcctgaa tgtgtccttc agcgtcagca 180  
accgtgtcgg gccctggtgg aacatcaaatt catgccagca gaagtg 226

&lt;210&gt; 25

&lt;211&gt; 265

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 85

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700268788F6

&lt;400&gt; 25

cggagggtgct cccaggcggc tgcactggct cggaggagta gcaggaggag ctccgcgcag 60  
gaacaaacct ggaggcaaac caganggagg caatgtttga atgactgtaa gaagaccaga 120  
cagtgaaaat gtcagccctc aactggaagc cctttgtgta cggagggtg cctccatcac 180  
cgcagaatgt ggtacatttc caattgattt gactaagact cggcttcaga ttcaaggcca 240  
gacaaatgat gccaaagtcc gagag 265

&lt;210&gt; 26

&lt;211&gt; 257

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 58, 99

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700270924F6

&lt;400&gt; 26

ctgggatccc cagggtcaat gggcatcctg ttcttgagc agggcactgt gagaaagnct 60  
ctcaccgtga ccaagtttct ctgagtgtcc agccaacna ggctcaccag ctccctccag 120  
ctaccgcccg tccatcaggt cagctgcca cccaggctg aacaccaacc ccagctatga 180  
gctcctggag gcatgactcc ctacgggcca gcagctccga tccctccca gtagttatca 240  
ttggcaatgg ccctcgg 257

&lt;210&gt; 27

&lt;211&gt; 244

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;



<221> unsure  
<222> 2, 8, 50, 56, 63, 76, 177, 219, 233, 240  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700303722F6

<400> 27  
gncgaggnc a ccaagggtgtt tctgcctcta cttagaaagt ccaaggggan gctggntaac 60  
gtnagcagca tggganccat gattccattt cagatgatgg ccgcctacgc ctgcacgaag 120  
gcagctataa gcatgtttctc agcccgatcat caggcaagag cttccaaatg gggagtnaaa 180  
gcgggaccat cattctggag cttcaaacca acatcgtag ctacacaggac agntgggatn 240  
aaag 244

<210> 28  
<211> 263  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700502047F6

<400> 28  
ggaaatgact ggtctgaagc ggcttggcag cctgagcagt cagggtactgc ggctctactg 60  
gcactgcctg aggggtccaag gactgtggtg attctcatgg aggaccctga gatttctgca 120  
atctgatcag tgtcaaagtc cactggattc gctctgagac tcttgcccta gaggatggcc 180  
aaagggctcc tgatgacctg tgccctttgg cttttggggc cctgttggac tacaccacct 240  
gtatctggga agggacagcc atc 263

<210> 29  
<211> 259  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 52, 137, 167, 189, 235, 254, 256  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700502260F6

<400> 29  
ggactacccc caccatgccc gtgtcattga gactctgatt gtccactatg gncgtgtctt 60  
tgaggaggag ccagaagaag cagctggcag ccaagagggg gcgtccgcca gtgtgcccag 120  
ctggagactg ctgaggncat tgtcttcccc cagcaggagg aggcggncga tggaaaccga 180  
gaatcccang tgcataatg actcagactc agagctggaa gaggtttctg acctntttcg 240  
cctcgagcgc cacnncctc 259

<210> 30  
<211> 260

<212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 140, 158, 161, 170-171, 177, 191, 193, 199, 203, 206-207, 212  
 <222> 217-218, 220, 224, 226, 231, 238, 243, 246, 248, 253, 255, 258  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700142213F6

<400> 30  
 cacaggggttt tccccaagag cacgcctctt tacttcaggg aattctcgga aactttcagg 60  
 attcggggggg cttccctggg aggagacgag agaggattag aagcggacat ccacggcttc 120  
 ttgtgatgac cacgcctttn gtctttgcta gaactctntg ngacttcccn nggtganttc 180  
 taatcacgga ntncacgcna agnttnngga antacgnngn cccntnaaga naaacacntt 240  
 ttngngnggg ggnanaantc 260

<210> 31  
 <211> 288  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700229555H1

<400> 31  
 caggagtacc actcacaggc cacctggcag gaagagataa gccccagcc cccgacatcc 60  
 aggacgcccc gaacctgcca atgtgtgtag ctatacctta ttacctcacc atgtgaaata 120  
 gccaatcata tgtgaacatg tctatgtgcc tcgtttgaat ccaccaatcc ctgtaactat 180  
 gcatctgctt ctgtacgect gcttctgctt cccaatccc tataaaagcc ccatgctgga 240  
 gctgctgggc gcgcaagtcc tcctaagaga ctgtgtgccc gcagtacc 288

<210> 32  
 <211> 258  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 53, 244  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700626839H1

<400> 32  
 ggggtgcgct ggagttgcgc atgcgccttc ccgccgcgca gggcaaagggt ggnggcgctc 60  
 tgggtgaatgg ttggttgcgc tgcaagagcg ttttctggct tttggtggcg aaggcggcct 120  
 ggccgcgagg tgcagctgct ggtgggcagg tgtactaatg tctacagact atgagcttcc 180  
 agaatctctg gagagagtag aaagtctctg atgttatggt acctttaatc gggttcatac 240

attnggggtgg cacagaat

258

&lt;210&gt; 33

&lt;211&gt; 268

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 7, 11, 40, 53, 150

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700627890H1

&lt;400&gt; 33

```

gtacaangag ngccgggggct tgggtctagt tggaggggan gcagtggcca gtncagggt 60
cagatgagag agttagccga gttaggggca gctactagga tgggggcagg aggagaagcg 120
gggctaacta taaagaagac tagatttcgn cacagtgggt atgtggaagg cagctttcaa 180
accgcccttg tcaaacaaca cagggccagc agccttcaag accaggctat ccctgccgtc 240
tgctggcatg ggggcacttg taccgtcc 268

```

&lt;210&gt; 34

&lt;211&gt; 299

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700059105H1

&lt;400&gt; 34

```

tcatacaaac ccaactgtgt ggccaaatcc agaggtgttt gacccttatt gatttgcacc 60
agagtcttcc cgacacagcc actcattcct gcccttctca ggaggagcaa ggaactgcat 120
tgggaaacag tttgccatga atgaactgaa ggtggccgtg gccctgaccc tgctccgctt 180
tgagctgctg ccagatccca ccaggatccc aatccccata ccaagactcg tggtgaagtc 240
caagaatggg atctacctgc gtctcaaaaa gctccaataa tcttgacagc acaagacag 299

```

&lt;210&gt; 35

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700059610H1

&lt;400&gt; 35

```

aaacaacctg actttcttgc gtgtgaggag tgccttttat gggaacagca tcactacaa 60
tatgtcctct gatggccgtt tgtcccgcgc ggctgccag attgtcatg agcacacaga 120
tggagtgtatc aaaatgagga aggtcagct gcagaatgag gaagagcttc agaaggccag 180
gaagaagagg cacttgatt tcctggacat cctgttgttt gccaaaatgg aggatgggaa 240
gagcttgtct gatgaggacc tgcgtgcaga ggtggacaca ttcattgttg agggatcatga 300

```

<210> 36  
 <211> 296  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700059630H1

<400> 36  
 ggggtttctct gtattttaccc ctacaagatc cctggatggt gtctctgggt tcttccaagg 60  
 ggccttcctg ctacagtctat ttctgggtgct gttcaaggca gtccaattct acttacgaag 120  
 gcaatggctg ctcaaggccc tcgagaagtt cccatccacg ccttcccact ggctttgggg 180  
 ccacgacctg aaggacagag aattccagca ggttcttacg tgggtagaga aattcccagg 240  
 tgcctgctta cagtgggtct cagggagcaa aacacgagtc ctgctctatg accctg 296

<210> 37  
 <211> 286  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 204  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700062959H1

<400> 37  
 ggcccatgga gcacacccag gctgtggact atgttaagaa gctgatgacc aagggccgct 60  
 actcactaga tgtgtggagt aggagctacc accctcccac ccctcgtcc ctgtaatcac 120  
 ctaacttctg ccgacctcca cctctgggtg ttctgcctg gcctggacac agggaggccc 180  
 agggactgac tcttggcctg agtngtgccc tcttgggccc ctaagcagag tccgggtccat 240  
 tgtatcaggc agcccagccc caaggcacat ggcaagaggg attgac 286

<210> 38  
 <211> 289  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700606459H1

<400> 38  
 ggtgagtccc gtgtggagaa aatatacaag taagaccgct acgtgcctgg cgactggaga 60  
 tgtgatgggg cacagcgac agagagccat aatggcctca tcgtacaggt ctgggacgct 120  
 cagcaacacc ccagcaggca ctgcactgtc tagtggacaa gctctgttag caggaagagc 180  
 ttctctgctg ctgtccaaga aaggctggtc aaggtccct accacctaca catactgtgg 240  
 ttggagaaat caaggttcct ggcaaaagag agtagcttca cgggaggca 289

<210> 39  
<211> 79  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 39, 48, 66, 68, 72  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700608882H1

<400> 39  
cttaacgctc ctgccacgcc gcctccgccc gtgcaatgnc tctgtagncg gcgatctacg 60  
tacgtntncc cngccccgt 79

<210> 40  
<211> 248  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 8, 21, 24, 35, 40, 52, 78, 90, 104, 137, 169, 209, 220  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700483988H1

<400> 40  
tttagttnc aatcatggagg nctntcaggt gaggnctan ctcccaaagc cngcgctgag 60  
tctacaactt ctctctaanag gtgttcgggn cggcggttggg gtcnctgcgg aggcgggctaa 120  
atcgcccgca gtttctncca tggttgcgcc cgtgtgttg cgcgctctnc gtaagaacaa 180  
gacccttcgc tatggagttc ccatgttgnt gctgggtgtn agtgggtcctt ttgggtcttcg 240  
cgaatttt 248

<210> 41  
<211> 352  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 28, 324, 337  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700501033H1

<400> 41  
ctctctgcct atgttctgag gttggagnct ttattacaga agctgggtgca gaaaggagca 60

```

attgagaaaag aagttgtgaa tcaggcccga ctagaccaag tcattgctgg ggcaatccac 120
aagtcagttc gaagagagct tggactgccga gaaggtagcc ctgccccagg cttattgcag 180
ttgctgacac tgataaaaaga taaggaggca gcagagggaag aggtccttct tcaggccgaa 240
ttagaaggac atttcacttg acccaagacc agcaaggctg tcatgagcag aacatgatgg 300
aggagctcat agaagtgatc agcncatccc ctttggnetg ccaagtaatt gc 352

```

```

<210> 42
<211> 233
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700137747H1

```

```

<400> 42
gtttctccat agcctcagac cccacatcag tatectcttg ctacttggag gagcacgtga 60
gcaaagaggc taaccatcta atcagcaagt tccagaagct gatggcagag gttggccact 120
tcgaaccagt caaccagggtg gtggaatcgg tggctaattg catcggagcc atgtgttttg 180
ggaagaactt cccaggaag agcgaggaga tgctcaacct cgtgaagagc agc 233

```

```

<210> 43
<211> 243
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 33
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700141708H1

```

```

<400> 43
tgggcagaaa ggaagccctg cagagcatca gangcccagc tagagggaca acacagagga 60
gtaatttgct gacagacctg cagggatgga cctgctttca gctctcacac tggaaacctg 120
ggctctcctg gcagtcgtcc tgggtgctct ctacggattt gggacccgca cacatggact 180
tttcaagaaa caggggattc ctgggcccga acctctgcct ttttttggca ctgtgctgaa 240
tta 243

```

```

<210> 44
<211> 295
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 286
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature

```

<223> Incyte template ID No: 700302454H1

<400> 44

```
gcggggccgtg ggtgatctgg tcggtaccgg agagcgcagg ttgtatcacc aacatggggg 60
actctcacga agacaccagt gccaccatgc ctgaggccgt ggctgaagaa gtgtctctat 120
tcagcacgac ggacatgggt ctgttttctc tcatecgtggg ggtcctgacc tactggttca 180
tcttttagaa gaagaaagaa gagataccgg agttcagcaa gatccaaaca acggccccac 240
ccgtcaaaaga gagcagcttc gtggaaaaga tgaagaaaac gggaangaac ttatc 295
```

<210> 45

<211> 286

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700304405H1

<400> 45

```
cggaagtga ccaaggcact gagcggcatc taatgcacct ggagttggac atctcagact 60
ccaagatcag gtatgaatct ggagatcacg tggctgtgta cccagccaat gactcagccc 120
tgggtcaacca gattggggag atcctgggag ctgacctgga tgtcatcatg tctctaaaca 180
atctcgatga ggagtcaaac aagaagcatc cgttcccctg cccaccacc taccgcacgg 240
ccctcaccta ctacctggac atactaacc cgccacgcac caatgt 286
```

<210> 46

<211> 311

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 299

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700306096H1

<400> 46

```
gataggaaaa taattttatt taggtttttt aaaaaagtta actttcacat ataaatttag 60
acttaaagat tacagtgtat attttccaaa aggagcggcc ctgaagggtg gccagacaag 120
ctcgccgagt gggcacaggg acactcgctc cagaaggagc tcagggtggaa gcgctttctt 180
taatctttcca cagtggccct tccctgttcc tcaccggggc tatgactggt aagaaaaccc 240
acaaccatca tttggggcaa cagcatctca ctagatggga ataagaacat gtctaggang 300
aaagcacaag c 311
```

<210> 47

<211> 307

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700325693H1

<400> 47

```
gtgccctcac gcagcttaat gtggcctttt cccgggagca ggcccacaag gtctatgtcc 60
agcaccttct gaagagagac aggggaacacc tgtggaagct gatccacgag ggcggtgcc 120
acatctatgt gtgcggggat gctcgaataa tggccaaaga tgtgcaaac acattctatg 180
acattgtggc tgagttcggg cccatggagc acaccagga tgtggactat gttaagaagc 240
tgatgaccaa gggccgctac tcactagatg tgtggagcta ggagcttacc aacctccac 300
ccctcgg                                     307
```

<210> 48

<211> 300

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 21, 49, 199, 226

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700059571H1

<400> 48

```
ctgtgctcct gaggcaagc naggcattcc tccaagacac tgcgggtcng agcagggact 60
gttcacgctg gtgccctgtg aactctggtg gaggtcagcc aacagctgct gtgtctgagt 120
tgctgagagg agagagaatg gcttgcaact agttttcttt ccacgtgcca agtctggagg 180
agctcgaga agttttgcng aaggggctaa aggacaactt tgctctgtc caggtctctg 240
tggtcgactg cccagattta acaaaggagc catttacttt cccgtaaaag gcattctgtg 300
```

<210> 49

<211> 314

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 13

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700060610H1

<400> 49

```
gcaagattga ggnggagaag gacaacctga agtctgagtt ccatctggag aacttggtg 60
tctgtgggtc taacttgttt acggcaggca ccgagacaac cagcaccacc ctgagattcg 120
ggctcctgct ccttatgaag tatccagagg tgcaagccaa agttcatgag gaacttgacc 180
gtgtgattgg acgccaccaa cccccagca tgaaggacaa gatgaagctg ccttataccg 240
atgctgtatt gcatgagatt caaagatata tcactctcct tccttcagat ctgccccatg 300
ctgtggtcca ggac                                     314
```

<210> 50



<211> 312  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 78  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700063128H1

<400> 50  
 cggtcggtac cggagagcgc aggttggtatc accaacaatgg gggactctca cgaagacacc 60  
 agtgccacca tgcctgangc cgtggctgaa gaagtgtctc tattcagcac gacggacatg 120  
 gttctgtttt ctctcatcgt gggggctcctg acctactggt tcatctttag aaagaagaaa 180  
 gaagagatac cggagttcag caagatccaa acaacggccc caccogtcaa agagagcagc 240  
 ttcgtggaaa agatgaagaa aacgggaagg aacattatcg tattctatgg ctcccagacg 300  
 ggaaccgctg ag 312

<210> 51  
 <211> 248  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 64-65  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700139656H1

<400> 51  
 caaaatacaa caaggagata aaagtcatac agtttgtgct gctggcttat tagctctgca 60  
 tggngaggg gccatggtaa gttgccaagg cattcagata taaatgtaat taagatgccca 120  
 tgtttgcttg cagtaatgaa gttataatca gaaactgcta aagtatgata aaaacagtga 180  
 ttgtttatgc acttatggaa gacaaagtga agtgatgtgg tttcttcaga acaggtgatg 240  
 cactgagg 248

<210> 52  
 <211> 115  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700141348H1

<400> 52  
 gtaaaagatg tggccagag aacgtaggaa catgcctgga gagacctaat gtgctcttgt 60  
 tctgcaaacc catgggcatt atttcctct cgcctcaaga gtcatactg gaagc 115

<210> 53  
<211> 249  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700182318H1

<400> 53  
agagtttctt tttgtccccc aacctgtagt tctaagttca acaaaacagt catcaacaaa 60  
agtgacggag gcttccatca gtgtcagggc tctgtgcaga cccagaatcc ctgttcccta 120  
tgtcatgttc cagcattgta tagcacgggt ccatgtcaca aacagaaagg tcaggaacac 180  
tgaggctctg gaatgtcact gctgcagcga ggcatgtca ctctctgtc tactctgtca 240  
gtgtcttac 249

<210> 54  
<211> 296  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 22  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700225376H1

<400> 54  
agcaagccta tttctgactg gnetgctgtg cagaatctag accactggca gtgggtgaca 60  
gccaggttga ggttaatcga agtctcgtcg caggetctgc tgtaagtctg gcctcttggc 120  
ctcacatctt ctttgtggga tcttcccta tctccagctt cctcagctgg tcagggagat 180  
ttggtccaga actagaagcc ttaataatct gagcaggtaa gagaggagta aaatgtacag 240  
tcttggacat tgactaaagg gtccctgcaga ggatatcaag gtaagtggct tggagg 296

<210> 55  
<211> 169  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700225757H1

<400> 55  
gctttctggg caagtctatg ttgcctagc tgaccagaa tttgctatat agactattct 60  
gtctcaaact cacagaaatt ctctgectg tgctctctga gcgagcacca ggattaaagg 120  
cgtgaatcgc tgtccccgtc ttttttctt cttcttttaa taaccact 169

<210> 56  
<211> 191  
<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 190

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700268788H1

<400> 56

```
cggaggtgct cccaggcggc tgcactggct cggaggagta gcaggaggag ctccgccgca 60
ggaacaaacc tggaggcaaa ccagaaggag gcaatgtttg aatgactgta agaagaccag 120
acagtgaaaa tgtcagccct caactggaag ccctttgtgt acggagggct ggcctccatc 180
aaccgcggan t                                     191
```

<210> 57

<211> 249

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 45, 118, 128, 163, 245

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700270924H1

<400> 57

```
tgggatcccc aggggctaata gggcatcctg ttcttgacgc aggggnactgt gagaaagtct 60
ctcaccgtga ccaagtttct ctgagtgtcc agccaaccca ggctcaccag ctccctcnag 120
ctaccgcncg tccatcaggt caactgcca cccagggtg aanaccaaac ccagctatga 180
gtcctctggag gcatgactcc ctgaggcca gcagctccga tccctcccag tagtgatcat 240
gggcnaggg                                     249
```

<210> 58

<211> 294

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 19, 32, 35, 104, 131, 188, 220

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700291661H1

<400> 58

```
actagcttct ccagctcctt ctttcccag angcncaggg gacctcggcc tccagcttac 60
cgggcggatc gaagcagcgg tcgggatggg actgctgggc ttgntgcagt caggcggctc 120
```

ggtgctcggg naggcgatgg agcaggtgac aggaggcaac ctgctttcca cgctgctcat 180  
cgcttgcnc cttcacgcta gccttgtcta cctgttccgn ctgcagtggt gccacatggt 240  
ccagctgccc gctggagcga aaagtcgcc atatatttac tctccaatc cgtc 294

<210> 59  
<211> 304  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700301979H1

<400> 59  
gatataat aa tcaaaaagaa aggaccacga ctcatgacct cccatcttcc catgcatctt 60  
ttctccaagt ctctcttcag ttccaaggcc aaggtgatct atctcgtcag aaatcccaga 120  
gatgttcttg tttctggtta ttatttctgg ggtaattcaa ctcttgcgaa gaagccagac 180  
tcaactgggaa cttatgttga atggttcctc aaaggaaatg ttctatatgg atcatgggtt 240  
gagcacatcc gtgcctggct gtccatgcaa gaatgggaca acttcttggt actgtactat 300  
gaag 304

<210> 60  
<211> 293  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700302770H1

<400> 60  
gtagccactc taactagggg cgtgctgaga caagaccacc tcattcctct gctgcttttc 60  
agacaggact gtctgcccga cccaccatga tccaggctgc actgttctt ggctgtatct 120  
tactgtcctc ggtgaccgcc tttccatgga agactcagga tgggtggcctg ccccatcagc 180  
cagctggcac agaaactgag cctacacaac tgctctacag caagagtcc cctccgacct 240  
ccagtacctg tcggaacctc ctaagcatgg cgccctgcc ccctgtagtc ctc 293

<210> 61  
<211> 174  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700303111H1

<400> 61  
caacagcaaa cgtgcgacca actcttcggc tagtgaatct ctaagccgcg aagagtgtt 60  
tgaagtagct ttaggtggaa gatgtcagaa agtaactcgg cagagggtag cgacagaagc 120  
gaggagcagg tgtctggtgc taaagtcac gccaggccc taaaaacgca agat 174

<210> 62  
<211> 273

<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 52, 139, 249  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700303390H1

<400> 62  
ctcagatggc aggcattggca tgcattgggat cttgttccct gagacaaagg cngatgcaga 60  
gggcatgtga ataaatcatg aggggccac agcaggccag caggccatag ctgacctcat 120  
tctggaagtg agagttgang agacccagc tgggacagaa aaggtaccac gcctataacc 180  
atggcctaac cgagggccag cagtggcagc ctccctgaaa gggacttcca gtccatccac 240  
aggcacgna gaaccagcaa gacatagcca gcc 273

<210> 63  
<211> 279  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700303722H1

<400> 63  
gtgactctga gtgtttgagc aggttaacttc taccttttga cctctatcgc aacagggtcca 60  
aaggttccaa aggagctggc aggacactca gacaagatcc actggcttca ggtgtgccta 120  
gtcttgaggt tcagaaagac ggaggcagct gaatgtgggt ctgaaccaac aacatctagc 180  
tacaagggga gccactcttc caccagcga ctgtgactgt tctcacaggt ctgaatttcc 240  
tgttggtatt cacaagatg ctttttattt ttaacttct 279

<210> 64  
<211> 275  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700306343H1

<400> 64  
gagaaaggcc accacctagc taggtgaggt gtgccagcat ggtcctgggg gtctcactgt 60  
cccagccct gggacgtgg ttccgccatg caatccctt cgctatcttc acgtgttac 120  
ttctttatat cagtgtatgg ctctccatg agtggccctt tgagttgcca gctcaaagaa 180  
ctcagcagtc cggcctgtgg gaactcaagc tctctctcc ttctccagcc ctcaaccttc 240  
tgcttcctgt cacctcaggt gttttacaag gctga 275

<210> 65  
<211> 294  
<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700306615H1

<400> 65

```
catccgtggg ctggctcacg ccattgcct gttcctggag tatacagaca caagctatga 60
ggacaagaag tacagcatgg gggatgctcc cgactatgac agaagccagt ggctgagtga 120
gaagttcaaa ctgggcctgg acttcccaaa tctgccctac ttaattgatg ggtcacacaa 180
gatcaccag agcaatgcc tctgcgcta ccttgcccg aagcacaacc tttgtgggga 240
gacagaggag gagaggattc gtgtggacgt tttggagaac caggctatgg acac 294
```

<210> 66

<211> 283

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 2

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700330359H1

<400> 66

```
gnggcccaat ggccctgag taaaagggtg tcaactgagag tccaaaggcc cgagtaggaa 60
catgcggctt agagccagt tctgaccca gaacgtccac tctgtacag gtagatgagg 120
agggtgttcgg gtgcccgag gcggtatccc gcctggcttt cgccctagcc tttctgcaac 180
gcatggacat gaagccgctg gtggtcctgg gactgccggc cccgacggcc ccttcgggct 240
gtctctcctt ctgggaagct aaggcacagc ttgtcagag ctg 283
```

<210> 67

<211> 263

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700368974H1

<400> 67

```
tccttagctg ggggcggggg gggcagctct ctactatctg aacgaattta atgtgggagt 60
catgccttat acaacacagg aaattaatgt gtgatcta gctgatcta tgacttatta 120
caatacagga atttaatgtg tgagccatgc cttcaaaaca tgtctagaat ttctggaatt 180
ggccggaagt caacagggat tgcttattta acctttcaaa tcactcattg tgactagggc 240
acatggtctt gcgcttgcta tga 263
```

<210> 68

<211> 269

<212> DNA

<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 129, 133, 153, 163, 165  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700373118H1

<400> 68  
gtttgaagca gggcgtaggg aaaagcggga ttaaagagta ctacctttat tagctccttc 60  
cctccagaat aggtgcaaat cctccccat gcccatcttc tgccacctgg ggtaaggatg 120  
tggcactgnc agnctgtcag cccactgact ttnagtcttc agntngcagt ctgggcaaat 180  
accagcgagc tctgttgaa caagaccagg ccttcagagc atctgaacca ctgtggcctt 240  
ctctcctcag ccttcactgt ggcttttgc 269

<210> 69  
<211> 288  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 159  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700375521H1

<400> 69  
gcagccatgg atcgcgggga ggaacctctg tccgcgaggg cggcgctgga gaccgagagc 60  
ctgcgattcc tgcacgtcac agtgggctcc ctgtgggcca gctatggctg gtacatcttc 120  
ttcagctgcy tctttctcta cattgtcatc cagaagctnt ccctgcgact gagggttcta 180  
aggcagaggg agctggacca agctgaggct gttctggagc ctgatgttgt tgtaagcga 240  
caagaggctt tagcagctgc tcgtttgaga atgcaggaag atctgaat 288

<210> 70  
<211> 280  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 16, 84, 96, 112-114, 118, 171, 226, 228, 236, 240, 267  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700461546H1

<400> 70  
tgaacaaact tctganaact tagttgacag tgttttgagt caactgaaaa aagcatgact 60  
tttggaaact ctgaatgcct tggntctcag tattancact ctattgaatt tnnntctnat 120  
taaagtatgt agtttttttag acttttttcc tgacagtatt atgtaatttt ntggcggtggg 180

tagatgggag tgtogettgt atgttaccat acagctgaca tgtatntntt gtctantctn 240  
attatcttag tagtttcctg ctgtggnatg taccataacc 280

<210> 71  
<211> 271  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 25, 64, 70, 82, 102, 155  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700480077H1

<400> 71  
ctgacctgac ccatgatgta agggncgta ggggagcatc accactgcaa aggctgacta 60  
aggncgtgtn ggctaaaggt cnccttgaag ccagtgctct anagtcacac cttctttgct 120  
ctgggcccag gaggcctact tcttcttttt ctcgnggaat cctggaatct taaagataaa 180  
agaacctaga aagaaaatca aaccacttt ccttgtgggg cagatggtaa tatgggactg 240  
agaacagcaa acctgggggc ttggagagga g 271

<210> 72  
<211> 210  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 187  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700480949H1

<400> 72  
gggcagaggt ccaggggaata agggaggctt ctaccaatga ttttgtttta tggtgcttga 60  
cagagatatt gtatgggttct ctgagagctc ccctgaaaac cttacctcca accacacaag 120  
ggttcctccc agagagcgct cgctgggcag caaggacaca ctcccatatt gccaaagcata 180  
tcaagtnccc aaagattggc agaaaattcg 210

<210> 73  
<211> 256  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 70  
<223> a or g or c or t, unknown, or other



&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700483259H1

&lt;400&gt; 73

gtgacgtaca tggaaaacaa agcctacggg gacaggctca agccgcagac agcagcaagt 60  
aaagcgctn cggccctgaa gcatggcagc tatcccttcc agcggctcgc tcgtggctac 120  
ccatgactac tatcggcgta agtagccct cgcagcccc gccagggct ggccagggc 180  
tctgtggctg accgcctcc cttccagg agtctgggc tcctcgcca gcaacagctc 240  
cggcggaagt gcagag 256

&lt;210&gt; 74

&lt;211&gt; 259

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 219

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700483475H1

&lt;400&gt; 74

ctctcgact tcggcaatgg ctggattccc accgtcatca cggccttgt ccttgcacc 60  
tcccaggccc aagctggatg gctacaacat gattatggc acctttctgt ctataagaaa 120  
tccatatgga accacattgt ccacaagttt gtcattggc acttaaaggg tgccctcgcc 180  
aactggtgga accatcgaca ttccagcac catgcgaanc caacatcttc cacaaggacc 240  
ccgacataaa gagcctgca 259

&lt;210&gt; 75

&lt;211&gt; 264

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700498995H1

&lt;400&gt; 75

gacagtagat gcccccaaag ctctagtaga tgatagtgtg ggggtgtgt gcggctccta 60  
cctgtgctgt tcattcacag tgcagtttaa gggagcaggc gccactgcat tccttggtg 120  
tgccctgagg gtgcttgctg ctttatatag taacagtcaa ttaaggtttc ttccagggaag 180  
agaaaaggga tggttttgag gggctcagaa aataggattc agtgtgtaac ataacaggta 240  
ggttgctggc acatgctgat atcc 264

&lt;210&gt; 76

&lt;211&gt; 271

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

<221> unsure  
 <222> 218, 228, 255, 270  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700504333H1

<400> 76  
 gtttatatttg acacagacat ggacaaagcg atggagcgct atgtctctat gcccaaggaa 60  
 aaggctccag aacacattcc ccttctcttc attgccttcc catcaagcaa ggatccaacc 120  
 tgggaggacc gattcccaga cccatccaca atgactgtgc tggtaaccac ggcctttgaa 180  
 tggttcgagg agtggcagga ggagcctaag ggcaagcnaa gtgttgcntt ggaaccctca 240  
 aaaaaacttc ccggaaccc tttatggggn a 271

<210> 77  
 <211> 167  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 11-12, 17, 21, 24, 48, 66, 72, 96, 128, 135, 162, 166  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700505040H1

<400> 77  
 caacaatctt nngtggncgt nctntctgga actgggcac atcagctnat gctgccatac 60  
 gctgntgga gngccgtggg gtgaaggctg cccgtncct ggtgggtacc ttcattgcag 120  
 cactaganat gcgtngtgtt tcccttactt tgatgcttgt gnatgna 167

<210> 78  
 <211> 267  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 5, 18, 26, 39, 90, 92, 122-123, 132, 137, 145, 152, 160, 168, 171  
 <222> 173-174, 186-187, 213  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700505423H1

<400> 78  
 ggggncttct gtgaggcnct gatacncatc gaggtctgna ttcagccagg ccacatgaag 60  
 cccaagatg ggtggctttt cctgtatgan tnagtacaga tatatccatg gccggggaat 120  
 tnngactggc anggtcncca ggganacca gncagctttn tcaagaantc ntnggttcc 180  
 cttggnntca caggaaccta ttaccttca tgnggtctgg ggttctggat ttaggtctt 240  
 tgggacagtc ccagttagaa gccttg 267

<210> 79  
 <211> 267  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 14, 20, 22, 24, 81, 248, 253  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700510534H1

<400> 79  
 tgaaagcgta aggncttgcn tntnagagge tctgtagtga gttctgtttg cctataaggg 60  
 aagtggaaca accgagacac ncgcaattct ttcgagtgtt aaggagcctg ggaggagcag 120  
 gcagccgctt gctttgagca tgctcaggtg ggtctgtccg ccgctgtggg gaaggcaccc 180  
 tgcagcaggg cttcctgccc caccctctcca ttgtagtagt gtccagatct cagaaacgca 240  
 gcttgaancc agnttcaaag gtaccag 267

<210> 80  
 <211> 291  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700513027H1

<400> 80  
 gggagggtttt aaaggccata ttgccaaacct caccgaaagg tttcaggaac ccgaggaagt 60  
 gttaatgtac aactcaccac ttcacgccac ccgaggetga agttgacgtt gccttttaag 120  
 cctttttaca tacactggcc atttcagaaa attctcaaca ataatgtctg ccttcgagtt 180  
 taagtcatgg tgtttttttag aattgacttg aaatgaaaat atcacaaagt gaatatatca 240  
 gctggtgatc gagtgactga aacccccctg gtctgcgggt gaccagttca g 291

<210> 81  
 <211> 273  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700528082H1

<400> 81  
 ataaaagtga aaactgggca agggcagggg gctgggcgtg aaccgcttac tagataatgt 60  
 tctctaaaaa ttggtcttga aaacctgtt tgtgtattcg ttttatgagt gcttaaaaaat 120  
 ggtgtgacca gggcatggc actgtcattg gaacagcaac atgcttgctg gtcacattgg 180  
 aatggggaaa tgtgaagaaa gctggacatc aggcctgagg caccatttc tttgtatgaa 240  
 agtgttgtgt acaaaccccc cactaatcat ttt 273

<210> 82  
<211> 268  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 68, 121, 165, 174, 182  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700528176H1

<400> 82  
caaggccctc agggaatcag aggagccagc ctgatccctg gtctctggag tcttaaaaca 60  
agtgtgtntt tgcaggtagt cctagttggg tgctgggggg aggctgccag gcctggcatg 120  
ngacattagg caggaagcca ctctggatga ttgtgcacat gaganctag tcanggaggg 180  
anggttttaa ggagaggact tagaatacaa gtgagaagcc agccgaggaa agggaaccaa 240  
gtcctcagaa tagaaggcta tactggct 268

<210> 83  
<211> 289  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700534427H1

<400> 83  
aaaacaggca agactttgga gaaagcagac caggtatgat ggccactttg ccaccaacag 60  
cccacacttc ccagcaacct gtaaacatag aggacgaaga tgggatcctg gatgagtatg 120  
accagtacag cctggcccaa tcttatgtcg tcgggtggagg tcggaaagga cgtaccaaga 180  
gagaagctgc tgccaacacc aaccgcccc ggcctgggtg gcatgagagg aagctgctga 240  
ccaagttcca gaactctgaa aggaaaaagg cctggcgctg agacagagc 289

<210> 84  
<211> 290  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 58, 157  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700535328H1

<400> 84  
ggtctgacca agtgagaaag acagcagggg caccagggc tcagacactc tggcgtaanc 60  
caaagaaag atggccacag ccagctccc tggtagcaag ctgtcatccc taaactctgc 120  
tctgtgcccc ttgtgggcag acgttaatca agccctngcc ctttctgatg ggccctccca 180

tcccgggaac actaaaaggt agtcttactg tccaccaccc tacacctgtt ttcataagtt 240  
atgcacaaat gcgaacagct gagacagaga tggagaagtt ctctgttttg 290

<210> 85  
<211> 275  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 17  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700607183H1

<400> 85  
cacagccct accagencac cctccataac tgcaccaaga ggatctatcc aacacctccc 60  
tgagcaggag gagcctgaag actccaaggg aaagagtcct gaggaaccct ttcctgtgca 120  
gctggatcta accacaaacc cacagggtga cacactggat gtctccttcc tctacctgga 180  
gcctgaggaa aagaaactgg tggctctgcc ttccctggg aaggaacagc gctccctga 240  
gtgcccgggg cccgaaaagc aaagaacccc ctgat 275

<210> 86  
<211> 285  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700607235H1

<400> 86  
ctgaagaccc accatgtctc tgctgactac tgtactactt ctctgggggtt tcattctggg 60  
cccagcaact gacacagcct gtatattcaa ggaagcctcg gaaaacagtc ccttgcccag 120  
gccctggctt tctgccaatc cagtgccctg gatcacacct ggctgagga cattcctgct 180  
gtgccagggg acagtgcggg atgtagtctt catgctgagg cgggaaggag atgatgggtt 240  
cctggcgata gtccaacaga tgtttttctg gagggagctg gaccc 285

<210> 87  
<211> 260  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 246-247  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700607396H1

&lt;400&gt; 87

```

ggccaccaag atggcgggcgc ccagcggacg gtgcgcgagg ttcgtcagct gacttgttct 60
cggagctgtg gccgcgaccc gcttctacct gtcccagtg accagagctc agtgaccagt 120
ccttatagtc gaaagcaggg tttttactgc tgaggacctg gacccgctgg gaggcttgcc 180
atggtaacag aacaggaggt agaggccata gggaaaaccc tagtggactc cacgcagccc 240
ctgcannccc gcttccgtgc                                     260

```

&lt;210&gt; 88

&lt;211&gt; 181

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 27, 180

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700607505H1

&lt;400&gt; 88

```

caaagaaaga aaacactcct ccgaggncgc cagcaaaggc agagaaagat ctgcaggatg 60
accttcatta cagcaggtgt tatattttat cttttttgcc ttcgtttcta gtgaatgtat 120
cactaaggtc ttcaaagaca tcagctttca aggaggtgcc taagtactgt ttccacacan 180
c                                                         181

```

&lt;210&gt; 89

&lt;211&gt; 280

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700607713H1

&lt;400&gt; 89

```

aattgagggg taaacatgtc tttgtgaaat atatgttctt ttacaatact ttgtactaat 60
ttacgtggaa ttattatttg tttctcattg gagatattta ttcagcctca atggcctttc 120
aggaactcct gaatcaggta ggaggcctag ggagattoca gacccctcag atgggttttg 180
ttgtcttcac cagtgttatt gtggtacctc atattataat agagaactta ctgcagccat 240
tcccagtcac cgctgctggg ttcctatcct cgacaatgtc                                     280

```

&lt;210&gt; 90

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 6, 77, 128, 143, 146

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

<221> misc\_feature

<223> Incyte template ID No: 700607873H1

<400> 90

```
gaaganacac caaagctcac tcaactcttca ggttctcact gaacctactg tateggaagg 60
gacttcacct cccagangtc cttttttatg aagactgttg agacagcttt ccagaaacta 120
gaaccatntg gaagatagac ctngngtat tectgtgcgg attatcttga ttacgttaat 180
taattctgga tgggactagg cttaaagtgc atcatgattt tccattaaca aggtgcacag 240
atgctacaaa tggctgggag aaatcct 267
```

<210> 91

<211> 258

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 11, 109

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700607972H1

<400> 91

```
gaggctaagc ntgtgcctcc tggctctctt ggggcagtcg cccgcgcgcc aagacctttc 60
gctgacctca gcgtcccgcct gctgcgcaag gaagggcggg gccactgcng tctggacagc 120
gtccgaaggc agcgagtcct ctggaggccg ccgtagtgca gaggagtggg ttgtcacgtg 180
acccaaggtt agaccatggc ttccaccaag ccgctgtctc gcttctggga gtggggcaag 240
aatatcgttt gcgtgggg 258
```

<210> 92

<211> 276

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700608519H1

<400> 92

```
caccacactg catctgcctt acatgccacc ttacaccatc atctatttcc cctccccagg 60
tgtggcctca gcgtttttcaa gatgaagggc acaagccata gtcactcctg gggaaccctt 120
caacactggt gctggaggga tgtggccaag accacgttgg gggcaagacc gagacttggg 180
gcgggactac aattgtggtt ggtggggcca ggactgacct cttagcctcc ataggcagct 240
acactgtctg tcacttttcc tcttccctg tgccgg 276
```

<210> 93

<211> 295

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 35

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700608661H1

<400> 93

```
cttagtccct gtactctgag ggtaagcctc atcgntcagg atctattgct gctgcttctc 60
cagctggctt ctgtagagac tgacagagac aactttgcag gataaggtag ctatcaagat 120
gctccatttg cgaagtacac agatgctgca gatgttggag agctccttaa ggaaatacct 180
tcctgagtc ttaaagggtt atgggactgt cttccacatg aaccagggag cccatttcaa 240
gctcaaggct ctggtggaca agtggcctga tttaatacag tggttgtccg tcctc 295
```

<210> 94

<211> 293

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 16, 179

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700609074H1

<400> 94

```
ggcgtggagt tggagnagag cgtcaggcgc ctccgggaga agtttcatgg aaaagtgtcc 60
cccaagaagg caggggctct tatgaggaag tttggcagcg accacactgg agttgggagc 120
tctatcgtgt acgggctcaa gcagaaagat ggacaggagc tgagcaacga tttggacgnc 180
caggaccac cagaggacat gaagcaggac caagatatcc aggcagtagc cacctctctg 240
ttgcccctga cgcaagccaa tcttcgaatg ttccaaagag cccaagatga cct 293
```

<210> 95

<211> 288

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 66, 138, 152, 236, 242, 252-253, 279

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700609967H1

<400> 95

```
ctgcctcgca gcccagagcg cgcgcctctc cagctccgc tccggttcc ccaaccaggc 60
ttattntggc tcccagaccg gtgcagaccc ctgaccggc ctccgcccag ctccgcaaaa 120
tgcgctactt tacttggnag gaggtggcgc angctccggg agggagaagg agcgatggct 180
cgtaatcgac cggaagggtg acaacatcag cgacttcagt tcgcccgcac ccggngggct 240
cncgggtcat cnnccactag ctggtcagga tgccacgna tcctttgt 288
```



<210> 96  
 <211> 164  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700625315H1

<400> 96  
 gtttagagcag ttacactgaa ccaaagtgaac tgagtttgta cagacggtaa tccgtaccaa 60  
 gcacactcac tgcctgatac tgaacaccca gcaagggtca tgcctgtgct aagtttgca 120  
 cattgtgttc ttttgcatc ttttttact tttattaaag gtgc 164

<210> 97  
 <211> 225  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 28, 59, 65, 73, 79, 84, 88  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700627089H1

<400> 97  
 cggggcggct gcagcaggcc accatgggag agcttcagga ggtgcagatc actagaggng 60  
 aagcnattgt tgcaggtnt ggcnttantg gttgagtcta tcaactccagt gggttccctg 120  
 tttgctctgg catcactc catcatctc ctcaagcttt tctcctaccg ggatgtcaat 180  
 ctgtggtgcc gccagcgaag ggtcaaggcc aaagctgtgt ctgca 225

<210> 98  
 <211> 265  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 62, 264  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700629293H1

<400> 98  
 atgaccttta acttttctaa aaatgtgaag tttgtactt atatatatca gctaaagtat 60  
 tntagcattc ttagtgtac ttagttgat gccactttta gtgtttttgt tgcttttgc 120  
 tgatttttat gaatgttcat ttttaagactc cttgttgaaa tgggacagtt tcgttctttg 180  
 ataagcccga gaagaggatt cccttgggtg ttgacctcct ctgcatgatg tgcccaagca 240  
 tctgaactgc aaccaaggcc tttnc 265

<210> 99  
<211> 95  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700643961H1

<400> 99  
agtataacca ggggccatct gaaagttggt ctctagccag ataagccact atgagcggta 60  
agcatcagtg ctaacgcaga aacttcctga gcagc 95

<210> 100  
<211> 307  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 8, 20, 21, 23, 47, 302  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700061625H1

<400> 100  
gggaaganac actggcttan nanttgggtc tgaggggaagc tctctgngtg gatacatatt 60  
tataggattt ggggtgcaacg aactgtgtga ctagttcagt tcaattcagg gagcaggcag 120  
aaaccagcag gtattagaag agatgttcta tatacagagt tctgaggcac tgcagattct 180  
gaagaattcc ctaagggaagc acctccctga gtccttaaag gttatgggac tgtctccaca 240  
tgaaccaggg aaacccattc aagctcaagg ctgtgggtgga caagtggctg atttaatact 300  
gntgtta 307

<210> 101  
<211> 97  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 48, 90  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700062809H1

<400> 101  
ccctgagagt ctccaaaatg ttctagcttg ttataagaac cccattanct cactaaagct 60  
tgcactggct gactcctgga cggggttccn gaatgtg 97

<210> 102  
<211> 214  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700139104H1

<400> 102  
gagatgttcc ctgtcatcga acagtatgga gacatthttgg taaaatactt gaggaagag 60  
aaaggcaaac ctgtccctgt gaaagaagtg tttggtgcct acagcatgga tgtgatcacc 120  
agcacatcat ttggagtga tttgtattcc ctcaacaacc cgaaggatcc ttttgtggag 180  
aaagccaaga agctcttaag aattgatttt ttgt 214

<210> 103  
<211> 265  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700139953H1

<400> 103  
aggaagccct gcagagcatc agaggcccag ctagaggggac aacacagagg agtaatttgc 60  
tgacagacct gcagggatgg acctgctttc agctctcaca ctggaaacct gggtcctcct 120  
ggcagtcgtc ctggtgctcc tctacggatt tgggacccgc acacatggac ttttcaagaa 180  
acaggggatt cctggggccca aacctctgcc tttttttggc actgtgctga attactatat 240  
gggtttatgg aaattcgatg tggag 265

<210> 104  
<211> 306  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 2, 61, 68, 111, 139, 263, (296)...(298), 305  
<221> unsure  
<222>  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700289281H1

<400> 104  
ancagatccg ggcactggag agggagctgc aagcaggagc agtcaagagt gtggtcagaa 60  
ncaccgtnag tggaccagca gggccggctt tttctaccat ggcggcccaa ngctatggct 120  
attaccgcac tgtcatatnc acagccatgt tggaggcta cagcctttac tacttcaacc 180  
gcaaaacctt ctcttttgtc atgccctcct tggtgatga gatcgctctg gacaaggacg 240  
atgtgggggc tcatcacgag canccagtcg gcagcctacg catcagcaag tttgtnnncg 300  
gggtnt 306

<210> 105  
<211> 183  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700303922H1

<400> 105  
ggcagcattg atccttatgt atatctgccc ttgggaaatg gacccaggaa ctgcattggc 60  
atgagggttg ctctcatgaa tatgaaactc gctctcacta aagttctgca aaacttctcc 120  
ttccagcctt gtaaggaaac acagatacct ctgaaattaa gcagacaagg acttcttcaa 180  
cca 183

<210> 106  
<211> 290  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 56  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700305783H1

<400> 106  
ctaacagtga atacatagct gcgatcctgg aactcagctc cctcatagtg aaacgncaac 60  
gccagccctt cctgtacctg gacttctgtg attgcctcac tgctgatggg cggcgcttcc 120  
gcaaggcctg cgacgtgggtg cacaacttca cagatgctgt catcagggag agacgcagca 180  
ccctcaatac ccaggggcgtt gatgaattcc taaaggccag ggctaagact aaaactttag 240  
actttattga tggtctcttg ctggccaagg atgagcatgg gaaggggctg 290

<210> 107  
<211> 177  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700329969H1

<400> 107  
gctatatcag gaggggaccc atgctgtgtc cttctgagat ctaacaggat taaccaatat 60  
gtaaaactaga ggaagtgggt ggctgcact gggcaagccc tctaggactc catccaagaa 120  
agaccagtgt gtgttgctct agaggcaaaag aaaccataa ggagctggca gtaaaac 177

<210> 108  
<211> 188  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 114, 116  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700368493H1

<400> 108  
aaagcttccc aatctgtgtg ctcaattggg aggatggcat taggagccag ggctggccat 60  
gggtacctta ctttcctccc tggggtatgc ccaggagaat ggagaaaaaa aaangnttta 120  
aagaaaaaat attttaaat tgatgctggc ctttttcaat tgtattgagt aaaagtgttc 180  
aagttgtc 188

<210> 109  
<211> 255  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 156, 157  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700376694H1

<400> 109  
ttatgattat caattttaca taacattaat attatatcaa acctccttaa gaaaatgagt 60  
atggatgttc acagtatgtt tgatttttat ctacaagaat gaatctgatt cagaatgctt 120  
ttcagctgac atacagagca ctaaataactt taagggnaac cataggtctg aatctcttaa 180  
gaattctcag tctctatggg atgtagggac gcattataaa tgcattaatc cttatagtca 240  
atcctgtgcc tagga 255

<210> 110  
<211> 284  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 16, 25, 54, 63, 68, 70, 80, 141, 154, 210, 275  
<221> unsure  
<222>  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No.: 700483803H1

<400> 110  
gaggaacgcc gccgcntcgc tcggnatcct acaccaatca ggaagctgct gtcnagccat 60  
gngggganen gagaagccan ctcaagaggc tgacgtggaa cccatggtaa catcaggggc 120

ctcagaagca gtgccaaggg ngctttcttg agancctcag aacatctctg atgtagatgc 180  
 cttcaacttg ctcttgagaga tgaaactgan acgacggcgt gaggtcccaa ccttccatgt 240  
 actgtgaccc agctagtggc cgaggatggc agcanggtgt atgt 284

<210> 111  
 <211> 258  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700502844H1

<400> 111  
 ggcagacctt agcagattct ggatgccatc agtgactgaa gccattaagg agactcggct 60  
 ggagtagcag ctaagaggac agagagacaa gggctacgag gcagcaatat aaacagatct 120  
 ggtgttgctg agatttgaga cgaaggtttc ccatggcttc ttttcacatc cgccagttcc 180  
 aggagaggga ctatgaacag gtctgtggata tgttctccag gggaatgaag gaacacatcc 240  
 ccaactgcctt ccgccact 258

<210> 112  
 <211> 250  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700503415H1

<400> 112  
 gtagctttcc ccttttgctg gcacagaagt ctgtccatct gcaagcgctt tggaacacag 60  
 actgcctgga gccaccttcc ttggggagac ctctctgcct cagctgtcgt cctgtgtcgt 120  
 cattcactaa agctcctgac gtcagattaa gcaagcagtg atgggttaca ttagagacaa 180  
 gccgcagaga taaggcctgt tgctgtttcg cagataatga tgagttttaa ttaccactg 240  
 gtttgtatgg 250

<210> 113  
 <211> 278  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 40, 44, (53)...(55), 60, 69, 72, 74, 159, 234  
 <221> unsure  
 <222>  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700514914H1

<400> 113  
 ccactgcagc gccccccccc aaagatggaa agattctgcn ttancttcat atnnncttan 60

aacatttgna cncnattctt taactttaga atctccccta gagcctgttc ttctttaaac 120  
accctttatg ctggagtagg atgatggetg agtttcttna aaagagctta aatatagagt 180  
cacaacatg agatagatgc ctgccgcca ctctttccac aaactegaga accnctttgt 240  
gcacgcagct gccatggaag gaaatcctgg ggcttctt 278

<210> 114

<211> 308

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No.: 700519169H2

<400> 114

cccaatcttt taagactttt cataatgatg ttaagaccag ggcagactat ctactggcca 60  
tggaactgcag ccaggagatg ccctgccct cctgggacct gcacaccacc tctctgggga 120  
acttgacaaa ggtccctaag gctaaggag gtctccttc tactaggctc ctgactttga 180  
ctctgtggtt ctctaggaac cgtgtgcaca cttgtctctg ttgtaaccac aaagggcagt 240  
agcacctagc atgtcatgtc ctgcccggc tgcttgctc ccaccaccc aggattctct 300  
gggctggc 308

<210> 115

<211> 124

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 34, 53, 87, 109, 122

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No.: 700535905H1

<400> 115

cctacctctg gtggaaaaag tacatcactc aggnccagct ggtccagttt gtngctgaca 60  
atcatccaga ccagctgcgg ggtcatntgg ccgtgctcct tccctctcng gtggctgtac 120  
tncc 124

<210> 116

<211> 262

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 148

<223> a or g or c or t, unknown, or other

WO 00/12760

PCT/US99/19768

<220>

<221> misc\_feature

<223> Incyte template ID No.: 700593984H1

<400> 116

```
gggaaagtgt gttctgaggg ccctgtgggg ccaaggggga ccagcctcac attccacacg 60
tgcgccactc tgcttgagc ctatttattt tgtatttatt tgaacagagt tatgtcctaa 120
ctatttttat agatttggtt aattaatncc ctgtcatttt caagttcatt ttttttattc 180
atatttatgt tcatggttga ttgtacctcc tgtcaccagc tgggtggggca ggggagacaa 240
ggtagaaagt ggccacagag tg 262
```

<210> 117

<211> 267

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 36, 37, 223

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No.: 700607844H1

<400> 117

```
ctcgatcgtc cagaccccac ggcgtcacca tgctgnncca tgatcaggaa ctactgttc 60
gggagcgtgg agacgtggcc ttggcaggtt ttaagcaccg ggggcaagga agacgtctcc 120
tatgaggaaa gagcctgcga agggggaaaag tttgctactg tggaagtgcac agacaaacct 180
gtggatgagg ctctccggga agcaatgcc aagatcatga agncgtgggg ggcaccaatg 240
acaaaggagt cggcatggga atgacag 267
```



## SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

CUNNINGHAM, Mary Jane

ZWEIGER, Gary B.

PANZER, Scott R.

SEILHNER, Jeffrey J.

<120> TOXICOLOGICAL RESPONSE MARKERS

<130> PA-0010 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/141,825; 09/172,711; 09/172,108

<151> 1998-08-28; 1998-10-13; 1998-10-13

<160> 117

<170> PERL Program

<210> 1

<211> 259

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700416862P7

<400> 1

```

gggggggga tttctggctc gagcatttgg agtttcttgg ctgctgaacg ggtagactaa 60
aggggggctg acatgggtga ggaggtacag aagcattctg tgcacacact agtggtcagg 120
tcattgaaga ggacccatga catgtttgtg gctgataatg gaaaaacctgt gaccttggat 180
gaaggagatc acaagcggaa aatggcaatc aagcttctga atgagtatgg cctgtgctg 240
catatgcta ctccaaaag                                     259

```

<210> 2

<211> 295

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 212, 227, 229, 232, 240, 243, 245, 250, 257, 269, 273, 277

<222> 288, 290

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700502957P6

<400> 2

```

gacctcttca ccatcgggac tagtactgc agggggcagc cctacctatt ccaactcaact 60

```

```

tgttacctot ggggtccag gcagggetta gteaaacctg cccagacacg gttaccttt 120
ttatgcccaa gctttcgggg tctgaggta ggggtgctt tctgcaccc ccaaggagca 160
gacactcag aatggagtca gctaggaacc cngggagctg cctcatnang cnettgatan 240
cangnacacn ttgcaactg cagacctntt tongaanaac attgcaangn teaac 295

```

```

<210> 3
<211> 273
<212> DNA
<213> Rattus norvegicus

<220>
<221> unsure
<222> 56, 220, 235, 237, 239, 249, 256
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700504502F5

```

```

<400> 3
tgacctgtct tctgtcaggt gactggtcaa gtgcagagcta gcttagttag tctggngtat 60
aaggcgccat catctctcca gtaagcctcc atcccaagc aactgaggct gtggcagtga 120
tgcacagcaac ctgtgtcacc caaaattatc cagcctcca cgggcactgc ctaggacctg 180
gggagggaag ggactttgca tcacatagcc tcaggttcgn gtttggtctt ggtangngnt 240
gcctgaaant ggtggnctcc agctggtgta cgg 273

```

```

<210> 4
<211> 264
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700505170F6

```

```

<400> 4
ggttggtctg cgatggcccg ctactctggc tctgtggaac catgtgtggg tccggcaett 60
gagactgga tctgaaagg ggtgaacctt cagcggaaac ttgcggcaa tttactccg 120
tcggacagc cccggcggga ggaggcagt aatgctttgt gctggggcac agggggcag 180
accagattt tggtaggatg tgcggacagg accgtgaggg accttaatgc ggaggagggt 240
acattccaga ccagagatad tgcc 264

```

```

<210> 5
<211> 268
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 231
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature

```

<223> Inocyte template ID No: 700511170F6

<400> 5

```
ggc*acaaga cgcgtgtgatt ggaagcaatg acgaagtcc caccctccgg ggagtgggta 60
tgtgtctgata catcatgtga tgggcagcct ggaggggatg cagggcggct ggagctatgt 120
ccaggggtggc atgggtgccc tctcagatgc cattgcaagc toggctactg cacatggagc 180
aagtatcttc acagagaaga ctgtggctaa ggtgcaagtg aacagcgaag nctgtccaa 240
gggttgtgtc tcagggcagg cgaaggat 268
```

<210> 6

<211> 384

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 28, 37, 44, 67, 71, 88, 102, 172, 238

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Inocyte template ID No: 700607827F7

<400> 6

```
gtgtacctac ctgttgagga aggaaggngt ggatggnaac cagnaacctg atgtccagca 60
caagggnaac nggcgtggct ttctaactta caagcctggg tnaactacagc ttgcagctac 120
ctaaccnatg caggaggcga aacctctgag cccagttgct attgtgacca tnaagatgtc 180
tkgcacacaa gcttccaccr agtctgggtt taatgggaag ttacctaacg attaccnca 240
gaagacacat gagacgcttg ctccaaagct ctgggatgca gcca 284
```

<210> 7

<211> 243

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Inocyte template ID No: 700500814F6

<400> 7

```
acttttttota atgtottatg gccacttctt atgagaatgg ggagctgtct tgcctgaggg 60
tgttgagagg aagcggccaga gcaggcccat catcccaacc cttggcctcg gcccttcccc 120
ctagctctg agcatttctt cagatctctt ttcttgagag tcaaggagac taaccacca 180
taaacagac acacctctg tggcccaaaa ggagaaaacy attagagggg tctctgtatg 240
atg 243
```

<210> 8

<211> 259

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 235

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700526819F6

<400> 8

```

ggcgagggcg gaaggggccc gtcacccgtcc gctgogacgt cggcgctgga gttgaacctg 60
gtgcgggctg ctttgccgtg tgagtogatg ggggcccga aaagagaacc ggaagagtgg 120
cgtctggaga agtatctggg tccctgggaag acatgctgca gccctgaaag tccaaagcaag 180
taaacctgccc cgggaagtgat cagtggatag tcccgcaaag tgactttctg aaggnatgct 240
gaggctgaga agtgactct                                     255

```

<210> 9

<211> 255

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 2, 4, 53, 55, 86, 88, 92, 106, 133, 211, 214

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700526176F6

<400> 9

```

cnangggccc caggggaatca agaggagcca gectgatccc tggcctcttg agnontaaaa 60
caagtgtgtt ttgcaggta gectangatg gntgtcgatg gagctncagc ctgcatggca 120
ttaggcagga agncaactct gatgattgtg cacatgagaa cctagtcagg gaggaggggt 180
ttaaggagag gcttagaata caagtggaga ncanogagaa agggaccaag tctcagaat 240
agaagctatc tgcct                                     255

```

<210> 10

<211> 269

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 30-31, 140

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700526082F6

<400> 10

```

ataaaagtga aaactgggca agggcgaagg nctggggcgt gaaccgctta ctagataaat 60
gtctctaaaa attggctctg aaaacccctg ttgtgtatct gttttatgag tgcctaaaaa 120
tggtgtgacc agggcatggn cactgtcatt ggaacagcaa catgcttget ggcacattgg 180
aatgggggaa tgtgaagaaa gctggcates ggcctggggc acccatttct ttgatgaaag 240
tgttgtgtca aacccccact aatcatttt                                     255

```

<210> 11  
 <211> 254  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 36, 47, 67, 80, 82-83, 92, 111  
 <223> a or g or c or t, unknown, or other

<320>  
 <321> misc\_feature  
 <323> Incyte template ID No: 700535328F6

<400> 11  
 caaggcgttc tgctgcgaga acgacattga catcgngcgc gtggtanacg tgcggaggct 60  
 ggcgccnate gtggcgcccn annacgagag gngcgccgcg tgagacctgc nttgcacct 120  
 cattogaac octaatgaag acacatggaa ggacccctgc ttggagaagc tcagtttgtt 180  
 ctgcgaggag agccgcagct tcaacgactg ggtcccgaga tcacctctcc gagtgcacgc 240  
 ctgcaggac attg 254

<210> 12  
 <211> 244  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 50, 102, 169, 171, 204, 210, 234, 242  
 <223> a or g or c or t, unknown, or other

<320>  
 <321> misc\_feature  
 <323> Incyte template ID No: 700368973F6

<400> 12  
 cacaattccc aaatggaaaa acttaaaaag gaattctgct gtgaaaggtn tatattactc 60  
 tagatttttc ttactgtaaa tattgtaxga ttgttaatact gncaatattt tattaccaa 120  
 caaatgttaa totatgtgaa atcagactta tttaaagggc tgctattang nggtgtggcc 180  
 ttgctgaca gattaaagtat attntgagtn agataactta ttaaggatgg aacnttaag 240  
 gntc 244

<210> 13  
 <211> 237  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 11, 20, 132, 136, 149, 151, 158, 171, 173, 184, 187, 190, 193  
 <223> 204-205, 208, 212-213, 215, 221, 227, 230-231, 236  
 <223> a or g or c or t, unknown, or other

<320>  
 <321> misc\_feature

<223> Incyte template ID No: 700368974F6

<400> 13

```
gtccttagct ngtggggcggg ggggtgcagt ctgtactat ctgaacgaat ttaatgtggg 60
agcatgcctt atadaadaca ggaattaat gtgtgatata atgggtgato tatgaattat 120
taccatagag anttangtgt gaacctgcnt ncaaaacngg tcagaatitt ngnaatggcc 180
ggantgnach ggntgattat taanntgnaa gngntggga naggecnggn ntgcgnt 237
```

<210> 14

<211> 335

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 38, 49, 53, 55, 107, 125, 132, 136, 138, 150-151, 163, 165-166

<222> 179-190, 184, 186, 189, 192, 199, 219-221, 223

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700369461F6

<400> 14

```
ctttaacggg tgggtgctg taagaatcgg tggcaggnet ctctctgeng gngntaatt 60
gtcttggaac gctactagga ccgaataact aaggccacat ctctacngtc taagagggga 120
aatangtag cnttgntncc acatgtggcn nagtgggggt gcngntatn gattaacann 180
tactantinc antgattant gtggtggtaa gatggctttn ntnaaaactg ccgcc 235
```

<210> 15

<211> 205

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 17, 19, 21, 56, 69, 75, 81, 103, 109, 111, 122, 127, 130, 137

<222> 148, 151, 153, 166, 174, 177, 190, 193, 195

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700782579F6

<400> 15

```
gggtcatitc caacctntna nacaagggga cgcacccaga catgccagt ttcacngaac 60
aagatgcang tccancagga ncagatagac tcagtcocat gantggctna ncaaccagcc 120
cngggcngcn caactgntgg acangtangg nantttctgc gacaantggt gantttnttg 180
cgaagctccn tgnacgaag aatgg 205
```

<210> 16

<211> 236

<212> DNA

<213> Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 11-12, 33-34, 46, 76, 85-86, 103, 123, 127, 144, 152, 157, 162

&lt;223&gt; 169, 174, 183-184, 186, 191-194, 217, 224, 230

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700782665F6

&lt;400&gt; 16

```

gggtctccatg xmgaccgggc gccttggggt tgmngagacg ctgcangccc ttaacgccc 60
cttgtagggc ccttgnaacc ccggnntggg ttaaggaaa cgnatgccc acaccttcgt 120
gtncganaact tttgggcacc gcgnaaactc gnttacngtg gnttttnang tagnggggtat 180
gtnnncgag nonncttagc ccggcttggt ctgcgggact gggtgcaan tgtctg 236

```

&lt;210&gt; 17

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700480927F6

&lt;400&gt; 17

```

ctggaccac atcacagaa atgaataaag cagatttctc tgttgagttc tgcagtaaac 60
cacttaaaag ccaatgtcaa gtcagccgca gacttaacta gcctgcctag cactgtagag 120
ggacttcaga agagtgtcgc ttccattggc aatacgttga acagtgtcag ccttgctgta 180
gaggcaatca agaagacogt ggtatgaacac aaggcacctt ggagttactg cagggcagtg 240
tggagacca tggaaagcaac caaatca 267

```

&lt;210&gt; 18

&lt;211&gt; 271

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 2, 92, 162, 181, 247-248, 253, 256, 262, 269

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700481732F6

&lt;400&gt; 18

```

gncttatltta tgtatgaaa tgcagaatc tgtacattcc tcaagccagt cctgtcgagc 60
caggtctgtc ccacccctgt acccaccacc antcccacct ggctgaaca ccccatgaga 120
cagagctgtg ctctgggctg gggcccccag gcctgggctg gncaggcaga cccatccccg 180
nagtcacatg gctccagtc ccgaggtctt cctgggctac aaagggggac cacacacacc 240
cagaatnntt tantgnattg gngggcceng g 271

```

&lt;210&gt; 19

<211> 283  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 17, 131, 199, 205-206, 208, 232-233, 239, 244-246, 251, 253-254  
 <222> 261-262, 266, 269, 274-275, 280  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700483143F6

<400> 19  
 cttaaattaa gatagaagtg aatgagacag atatctngta agaacctgta tttctttgtg 60  
 tgatcagatc tagtgtgggtg ggatgataga agttgaactt gctttattgc tatgggttaa 120  
 aatattttgt ntcatataaa tggcctattg aatgctttt ctgttctat aataaataa 180  
 cctgatgaaa aagtaaaaaa aaannnana aaaaaaaggg ggggcccccg cnnagggent 240  
 tttnncccg ngattantt nngccngnc cctnngggg cca 283

<210> 20  
 <211> 256  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700484538F6

<400> 20  
 ctggcctcag ctctctaagt totggggagtc ggacaggtgt ttgcacacac catggcccac 60  
 cggggaccta gaacctacag tgaacctga cccaggctct gtggatgttc tgcctcctga 120  
 ggtagacagc ctctaataac ctgttaggga cctaggacca gagctggggt gccaggcat 180  
 gtcccaacat gtgcctcagg ccacagggat atcggttgaa gtgcatttgg aagtgtgctg 240  
 ggacgcagc cagctt 256

<210> 21  
 <211> 272  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 64, 68, 124, 140, 145, 213, 260, 267, 269  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700480077F6

<400> 21  
 ctgacctgac ccatgattta aggacogtag tttagcaagg accactgcaa aggggggcta 60  
 agnctgnbg ggctaaaggt ctctttgagc ccagtggcta tagtcacacc ttctttgctc 120  
 tggncagga ggcctacttn tctntactc gtggactcct ggaatcttaa agataaaaga 180



acctagaaag aaaatcaaac ccacttttct tgggggaga tggtaatatg ggaactgagac 240  
agcaaacctg gggcttggan aggaacnanc tc 272

<210> 22  
<211> 270  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 40, 44, 46, 114  
<223> a or g or c or t, unknown, or other  
  
<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700060207F7

<400> 22  
taccocctgca tgggataccg tttctcgacc ccagtgcaacn tggntnctgt catcccatga 60  
aattgcagea agggcagctc cttctgtggg aacagattaa ctctacaca tggngtagat 120  
tcacacacctg ccaggaaagc agaagcatta cttaagtgtc ctgtgaaggc aaacatcaag 180  
tcaattcagc ttatcttgaa gagtggcaaa ccataaactc caaatgtcat tgtgtgaaac 240  
tgaacgatgg taatttcatt ccggtgctgg 270

<210> 23  
<211> 250  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 45, 49-50  
<223> a or g or c or t, unknown, or other  
  
<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700133105F6

<400> 33  
ccctcctgta tctgaaccca gttctcagc totgagatga gtsenggann ggttcccaa 60  
cctatgctca ataccacagg cagcctgcag gagggagaaa tgggtaaaat gttccatggg 120  
aaatgtctca gaatcgtctc cccgaatct cctgctaagc ttactgtctg ctatggagtg 180  
atcatggtcc tcagtctagc tctagtctct cttctgttg ctttgtcagt aaaaatgaca 240  
ccacagatct 250

<210> 24  
<211> 226  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700138117F6

&lt;400&gt; 24

gaggattcac tccatttgc tcccgctgg ccattgagtga gctgcccctt ctgagtcacag 60  
 agggagccag agggcctcac aacacacagag ggtctcagag ctccctggag gaaggctcag 120  
 ttacaggctc agaggctcgg caccagcttag gtgtccctgaa tgtgtccctc agcgtcagca 180  
 accgtgtcgg gcccgtgtgg aacatcacat catgccagca gaagtg 226

&lt;210&gt; 25

&lt;211&gt; 265

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 85

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 7002677886

&lt;400&gt; 25

cggaggctgct cccaggcggc tgcactggct cggaggagta gcaggaggag ctccgcgcag 60  
 gaacaaaccl ggaggcaaac cagangggagg caatgtttga atgactgtan gaagaccga 120  
 cagtgaacat gtcagccctc aactgggaag cctttgtgta cggagggtg cctccatcac 180  
 cgcagaatgt ggtacatttc caattgattt gactaagac cggcttcga ttcaggcca 240  
 gacaaatgat gccaaattcc gagag 265

&lt;210&gt; 26

&lt;211&gt; 257

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 58, 99

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 760270924F6

&lt;400&gt; 26

ctgggatccc cagggttaat gggaatcctg ttcttgacag agggcaactgt gagaaagct 60  
 ctcccgctga ccaagtttct ctgagtgtcc agccaaacma ggtcaccag ctccctccag 120  
 ctacogcccg tccatcaggt cagctgccaa ccccaggctg aacacaaac ccagctatga 180  
 gctcctggag gcatgactcc ctccgggcca gcagctccga tccctccca gtagttatca 240  
 ttggcaatgg cctcgg 257

&lt;210&gt; 27

&lt;211&gt; 244

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

<231> unsure  
 <222> 2, 4, 50, 56, 63, 76, 177, 219, 233, 240  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700303722F6

<400> 37  
 gncgagggnc ccaagggtgtt totgocctta cttagaaagt ccaagggggn gctggntaac 60  
 gtnagcagca tgggandcaat gattccattt cagatgatgg ccgctacgc ctgcacgaag 120  
 gcagctataa gcattgttct agcccgatc caggcaagag cttcraastg gggagtnaas 180  
 gcgggaaccat cattctggag ctccaaacca acatcgtang ctcacaggac agntgggato 240  
 aaag 244

<210> 28  
 <211> 363  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700502047F6

<400> 38  
 ggaaatgact ggtctgaagc ggcttggcag cctgagcagt caggtaactgc ggcctctactg 60  
 gcaactgcctg agggccaag gactgtggcg attctcatgg aggaccctga gattctctga 120  
 atctgatcag tgcacaatgc caatggatc gctctgagac tctgcctta gaggatggcc 180  
 aaagggtcc tgatgaacta tgccctttgg ctcttggggc cctgttggac tacaccacct 240  
 gtatctggga agggacagcc atc 253

<210> 29  
 <211> 259  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 52, 137, 167, 189, 235, 254, 256  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700502260F6

<400> 39  
 ggaactacccc caccatgccc gtgtcattga gactctgatt gtccactatg gnetgggtctt 60  
 ctaggaggag ccagaagaag cagctggcag ccaagagggg gcgtccgcca gtgtgcccag 120  
 ctggagactg ctgaggncat tgtcttccc cagcaggagg aggcggncga tggaaaccca 180  
 gaatcccang tgcattcaatg actcagaatc agagctggaa gaggcttcag acctntttcg 240  
 cctgggaagc cactncctc 259

<210> 30  
 <211> 260

<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 140, 158, 161, 170-171, 177, 191, 193, 199, 203, 206-207, 212  
<222> 217-218, 220, 224, 226, 231, 238, 243, 246, 248, 253, 255, 258  
<223> a or g or c or t, unknown, or other

<230>  
<221> misc\_feature  
<223> Incyte template ID No: 700142213F6

<400> 30  
cacagggttt tccccaaagg caagcctctt tacttcaggg aattctcgga aactttcagg 60  
attcgggggg cttccctggg aggagacgag agaggattag aagcggacat ccacggcttc 120  
ttgtgatgac caagcctctt gtctttgcta gaactctntg ngacttccn nggtgantt 180  
taatcacgga ntncacgna agnttmgga antacgngn ccctnaaga nnaacacntt 240  
ttnggngngg ggnanaaact 260

<210> 31  
<211> 288  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700229555H1

<400> 31  
caggagtacc actcacagga caactggcag gaagagataa gccccagcc ccgacatcc 60  
aggaagcccc gaacctgcca atgtgtgtag ctatacctta ttacctcacc atgtgaaata 120  
gccaatcata tgtgaacatg tctatgtgcc tggtttgaat ccacccaatcc ctgtaaatat 180  
gcactctgct ctgtacgctt gcttctgctt ccccaatccc tataaaagcc ccgtgctgga 240  
gtgtctgggc ggcgaagtc tctaagaga ctgtgtgccc gcagtacc 288

<210> 32  
<211> 258  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 53, 244  
<223> a or g or c or t, unknown, or other

<230>  
<221> misc\_feature  
<223> Incyte template ID No: 700626839H1

<400> 32  
gggtgcgagt ggagttggc atgagccttc ccgcccgcga gggcgaaggt ggnggcgctc 60  
tggtgaatgg ttggttgctg tgcaagagcg tttcttggtt ttggtggcg aagsgcgctt 120  
ggcgcgcagg tgcagctgct ggtgggcagg tgtaactaat tctacagact atgagcttct 180  
agaatctctg gagagagtac aaagttctgc atgttatggt acctttaatc gggttcatat 240

actnnggggtgg cacagaat

258

&lt;210&gt; 33

&lt;311&gt; 288

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 7, 11, 40, 53, 150

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700627830H1

&lt;400&gt; 33

```

gtacaangag ngccgggggt tgggtctagt tggaggggan gcagtggcca gtncagggct 60
cagatgagag agttagccga gttaggggca gctactagga tgggggcagg aggagaagcg 120
gggctaacta taaagaagac tagatttcgn cacagtgggt atgtggaagg cagctttcaa 180
accgcccttg tcaacaaca cagggccagg agccttcaag accagggctat cctgcccgc 240
tgctggcatg ggggcacttg taccgtcc

```

268

&lt;210&gt; 34

&lt;311&gt; 399

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700059105H1

&lt;400&gt; 34

```

tcatacaaac ccaactgtgt ggcacaatcc agaggtgttt gacccttacc gatttgcacc 60
agagttcttc cgaacagcc actcatctct gcccttctca ggaggagcaa ggaactgcac 120
tgggaacacg ttggccatga atgaactgaa ggtggccgtg gccctgaccc tgcctcgttt 180
tgagctgctc ccagatccca ccaggatccc aatccccata ccaagactcg tgttgagtc 240
cagaatggg atctacctgc gtctcaaaa gctccaataa cttgacagc scaagacag 299

```

&lt;210&gt; 35

&lt;311&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700059610H1

&lt;400&gt; 35

```

aaacaacctg actttcttgc gtgtgaggag tgccttttat gggaaacagca tcattacaa 60
tatgtcctct gatggccgtt tgcctccgag gccctgcag attgctcatg agcacacaga 120
tggagtgtac aaaaatgagga aggtccagct gcagaatgag gaagagcttc agaaggccag 180
gaagaagagg cacttggaat tcttggaat cctgttgttt gcaaaaatgg aggatgggaa 240
gagcttgctc gctgaggacc tgcgtgcaga ggtggacaca ttcattgttg agggctcatg 300

```

<210> 36  
 <211> 256  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700059630H1

<400> 36  
 ggggtttctct gtattttaccc ctacaagatc cctggatggt gtctctgggt tcttccaagg 60  
 ggccttctctg ctcaagtctat ttctgggtgct gtccaaggca gtccaattct acttaaggag 120  
 gcaatggctg ctcaaggccc tggagaagtt cccatccacg ccttcccact ggctttgggg 180  
 caacgacctg aaggacagag aattccagca ggttcttaag tgggtagaga aattccagg 240  
 tgcctgctta cagtggctct caggagagca aacacgagtc ctgctctatg accctg 296

<210> 37  
 <211> 286  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> ungure  
 <222> 204  
 <223> A or G or C or T, unknown, or other

<230>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700062959H1

<400> 37  
 gggccatgga gacaccccag gctgtggact atgttaagaa gctgatgacc aaggggccgt 60  
 actcactaga tgtgtggagt aggagctacc accctccacc cctcagctcc ctgtaatcac 120  
 ctaactttct cggacctcca cctctgggtg ttcttgcttg gctgggacac agggaggccc 180  
 agggactgac tcttgacctg agtngtgccc tcttgggccc ctaagcagag tccggtccat 240  
 tctatcagga agcccagccc caaggcacat ggcaagaggg attgac 286

<210> 38  
 <211> 289  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700606459H1

<400> 38  
 ggtgagtcct gtgtggagaa aatatacaag taagaccgct acgtgcctgg cgactggaga 60  
 tgtgatgggg cccagcgcac agagagccat aatggcctca tegtacaggt ctgggaogct 120  
 cagcaacacc ccagcaggca ctgcaactgtc tagtggacaa gctctgttag cagggaagagc 180  
 ttctctggct ctgtccaaga aaggctggtc aaggctccct accacctaca catactgttg 240  
 ttggagaaat caaggctcct ggcaaaagag agtagcttca cgggaggca 289

⑥ (Cont'n)

WO 0012760

PCT/US99/19768

<210> 39  
<211> 79  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 39, 48, 66, 68, 72  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700608882H1

<400> 39  
cttaacgctc ctgccacgcc gcttcggccc gtgcacgnc tctgtagncc gcyatctacg 60  
taegtntncc cngcccgt 79

<210> 40  
<211> 248  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 8, 21, 24, 35, 40, 52, 78, 90, 104, 137, 169, 209, 220  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700483988H1

<400> 40  
tttagtttca atcatggagg nctntcaggt gaggnectan ctccccaaagc cngcyctgag 60  
tctacaactt ctcttaanag gtgttcgggn cggcggttggg gtccctgcgg aggcggctaa 120  
atcgccgcga gtttctncca tgggtgcgcc cgtgtgtgtg cgcgctctnc gtaagaacaa 180  
gacctttgcg tatggagttc ccatgttgnt gctgggttgn agtggttctt ttggtcttgc 240  
cgaatttt 248

<210> 41  
<211> 352  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 28, 324, 337  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700501033H1

<400> 41  
ctctctgctt atgttctgag gttggagnot ttattacaga agctggtgca gaaaggagca 60

```

Attgagaaag aagttgtgaa tcaggcccca ctagaccag tcatigctgg ggcattccac 120
aagtcagttc gaagagagct tggactgcca gaaggtagcc ctgccccagg cttattgcag 180
ttgctgacac tgataaaaga taaggaggca gcagagggaag aggtccttc ttaggccgaa 240
ctagaaggac atttcacttg acccaaggac agcaaggctg tcatgagcag aacatgatgg 300
aggagctcat agaagtgatc agncatccc ctttggmctg ccaagtaatt gc 352

```

&lt;210&gt; 42

&lt;211&gt; 333

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700137747H1

&lt;400&gt; 42

```

gtttctccat agcctcagac cccacatcag tctctctctg ctacttggag gagcacgtga 60
gcaagagggc taacctctc atcagcaagt tccagaagct gatggcagag gttggccact 120
tcgaaccagt caaccaggtg gtgggaatcgg tggctaattg cctcggagac atgtgttttg 180
ggaagaactt ccccaggag agcagaggaga tgctcaacct cgtgaagagc agc 233

```

&lt;210&gt; 43

&lt;211&gt; 343

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 33

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700141706H1

&lt;400&gt; 43

```

tgggcagaaa ggaagccctg cagagcatca gmgcccaga tagagggaca acacagagga 60
gtaatttggc gacagacctg cagggatgga cctgctttca gctctcacac tggaaacctg 120
ggtcctcctg gcagtcgtcc tggctgctct ctacggattt gggacccgca cacatggact 180
tttaagaaaa caggggatcc ctgggcccac acctctgctt ttttttggca ctgtgctgaa 240
tta 243

```

&lt;210&gt; 44

&lt;211&gt; 295

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 286

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature



<223> Incyte template ID No: 700302454H1

<400> 44

```

ggggggcctg ggtgatctgg tgggtacgg agagcgagg ttgtatcacc aacatggggg 60
actctcaoga agacaccagt gccaccatgc ctgaggccgt ggctgaagaa gtgtctctat 120
tcagcacgac ggacatgggt ctgtttttct tcctcgtggg ggtccctgacc tactggttca 180
tcttttagaaa gaagaaagaa gagataccgg agttcagcaa gatccaaaca acggcccccac 240
ccgtcxaaga gaggagcttc gtggaaaaga tgaagaaaac ggggaangaac ttatc 295

```

<210> 45

<211> 286

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700304405H1

<400> 45

```

gggaagtga ccaaggcact gagcggcato taatgcacct ggagttggac atctcagact 60
ccaagatcag gtatgaatct ggagatcag tggctgtgta cccagccaat gactcagccc 120
tggtcaccca gattggggag atctggggag ctgacctgga tgtcatcatg tctctaaaca 180
atctcgatga ggagtcacac aagaagcctc cgttccctg cccacacacc taacgcaagg 240
ccctcactta ctactggac atcactaac ccgcaagcac caatgt 286

```

<210> 46

<211> 311

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 299

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700306096H1

<400> 46

```

gataggaaaa taattttatt taggtttttt aaaaaagtta accttcacat ataasttttag 60
acttaagat tacagtgtat attttccaaa aggagcgccc ctgaagggtg gccagacaag 120
ctcgccgagt gggcacaggg acactogctc cagaaggagc tcagggtggaa gcgctttctt 180
taatottcca cagtggccct tccctgttcc tcaccgggac tatgactggg aagaaaaccc 240
acacccatca tttggggcaa cagcatctca ctagatggga ataagaacat gtctaggang 300
aagcacaag c 311

```

<210> 47

<211> 307

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700325693H1

<400> 47

```
gtgcctctac gcagcttsak gtggcctttt ccggggagca ggcccacaag gtctatgtcc 60
agcaccttct gaagagagac agggaaacac tgttgaagct gatccacgag ggoggtgccc 120
acatctatgt gtgcggggat gctcgaaata tggccaaaga tgtgcaaac acattctatg 180
acattgtggc tgagttcggg cccatggagc acacccaggc tgtggactat gttaagaagc 240
tgatgaccaa gggcgcctac tccatagatg tgtggagcta ggagcttacc aacctccac 300
cctcgg                                     307
```

<210> 48

<211> 300

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 21, 49, 199, 226

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700059571H1

<400> 48

```
ctgtgctcct gagtgcgaag naggcattcc tccaagacac tgcgggtcng agcagggact 60
gttcacgctg gtgcctctgt aactctggtg gaggtcagcc aacagctgct gtgtctgagt 120
tgctgagagg agagagagat gcttgcaact agttttcttt ccacgtgccs agtctggagg 180
agctgcaga agttttgong aaggggctaa aggcacactt tgcctctgtc caggtctctg 240
tggtcgactg cccagattta acaaaggagc catttacttt ccgtaaaag gcctctgttg 300
```

<210> 49

<211> 314

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 13

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700060610H1

<400> 49

```
gcaagattga ggnggagaag gacaacctga agtctgagtt ccatctggag aacttggctg 60
ctgttgggtc taacttggtt acggcaggca ccgagacaa cagcaccacc ctgagattcg 120
ggctcctgct ccttatgaag tatccagagg tgcagacaa agttcatgag gaacttgacc 180
gtgtgattgg aogccaccaa ccccccagca tgaaggacaa gatgaagctg ccttatcccg 240
atgctgtatt gcatgagatt caaagatata tcaattctct tccctccagt ctgcccctg 300
ctgtggtcca ggac                                     314
```

<210> 50

<211> 312  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 78  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700063128H1

<400> 50  
 cggtcgggtac cggagagcgc aggttgkato accaacatgg gggactotca ogaagacacc 60  
 agtgcaccca tgccctganc cgtggctgaa gaagtgtctc tattcagcac gacggacatg 120  
 gttctgtttt ctctcatagt gggggtctctg acctactggg tcatctttag aaagaagaaa 180  
 gaagagatad cggaggttcag caagatccaa acsacggccr cccccgtcaa agagagcagc 240  
 ttctgtggaaa agatgaagaa aacgggaagg aacattatcg tattctatgg ctcccagacg 300  
 ggacccgctg ag 312

<210> 51  
 <211> 248  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 64-65  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700139656H1

<400> 51  
 caaaatacas caaggagata aaagtcatac agtttgtgct gotggcttat tagctctgca 60  
 tgganagagg gccatggtaa gttgccasgg catccagata taaatgtant taagatgcca 120  
 tgtttgcttg cagtaatgaa gttataatca gaactgcta aagtatgata aaascagtga 180  
 ttgtttatgc acttatggaa gacaagtgta agtgatgttg tttcttcaga acaggtgatg 240  
 cactgagg 248

<210> 52  
 <211> 115  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700141348H1

<400> 52  
 gtasaagatg tggteccagas aacgtaggaa catgcectgga gagacctaatt gtgctcttgt 60  
 totgcaaacg catgggcatt attccctctt cagctcaaga gctcatactg gaagc 115

<210> 53  
 <211> 249  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700182318H1

<400> 53  
 agagtttctt ttgctcccc aacctgtagt tataagttca acaaaacagt catcaacaaa 60  
 agtgacggag gottccatca gtgtcagggc tctgtgcaga cccagaatcc ctgttcctta 120  
 tgtcatgttc cagcattgta tagcaagggt ccatgtcaca aacagaaagg tcaggaacac 180  
 tgaggtctgt gaatgtcact gctgcagcga ggtcatgtca ctctctgtc tactctgtca 240  
 gtgtcttac 249

<210> 54  
 <211> 296  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 22  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700225376H1

<400> 54  
 agcaagccta tttctgactg gnetgtgtg cagaatctag accactggca gtgggtgaca 60  
 gccacagbga gyttaatoga agtctctgtg caggctctgc tgtaagtctg gctcttggc 120  
 ctacatctt ctttgtggga tcttcccta tctccagctt cctcagctgg tcaggggagat 180  
 ttggtccaga actagaagcc ttaataatct ggcaggtaa gagaggagta aaatgtacag 240  
 tcttggacat tgactaaagg gtctgcaga ggatatcaag gtaagtggct tggagg 296

<210> 55  
 <211> 169  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700225757H1

<400> 55  
 gctttctggg caagtctatg ttgccttagc tgacccagaa ttgtctatat agactattct 60  
 gtctcaaac ctacagaaatt ctctgctg tgctctctga ggcagcaca ggattaaagg 120  
 cgtgaatgc tgtcccgctc tttttcttt cttctttta taaacct 169

<210> 56  
 <211> 191  
 <212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 190

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700268788X1

<400> 56

```

cggaggtgct cccagggggc tgcactggct cggaggagta gcaggaggag ctccgcgcga 60
ggaacaaacc tggaggcaaa ccagaaggag gcaatggttg aatgactgta agaagaccag 120
acagtgaasa tgtcagccct caactggaag ccttttgtgt acggagggct ggcctccatc 180
aacccggan t                                     191

```

<210> 57

<211> 249

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 45, 118, 128, 163, 245

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700270924H1

<400> 57

```

tggyatcccc aggggctaatt gggcactctg ttcttgccgc agggnaactgt gagaaagtct 60
ctcacctga ccaagtttct ctgagtgtcc agccaaacca ggctcaccag ctccctcnag 120
ctaccgcneg tccatcaggt caactgccaa ccccaggctg asnaccaaac ccagctatga 180
gtccttgga gcactgactc ctccaggcca gcagctccga tccctcccag tagtgatcat 240
gggcnaggg                                     249

```

<210> 58

<211> 294

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 19, 32, 35, 104, 131, 188, 220

<223> a or g or c or t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700291661K1

<400> 58

```

actagattct ccagctccnt ctttcccgag angcngcagg gacctcgcc tccagcttac 60
cgggcccgat gaagcagcgg tgggatggg actgctgggc ttgntgcagt caggcggctc 120

```

```

ggtgctcggg naggogatgg agcagggtgac agsagggaac ctgccttcca cgtgctcat 180
cgcctgcnc cttcacgttta gctttgtcta cctgttccga ctgcagtggt gccacatggt 240
ccagctgccn gctggagaga aaagtcogcc atatatttac tctccaatcc cgtc 299

```

```

<210> 59
<211> 304
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700301979H1

```

```

<400> 59
gatatattaa tcaaaaagaa aggaccaaga ctcatgaact cccatcttcc catgcatctt 60
ttctccaagt ctctcttcag ttccaaaggcc aaggtgatct ctctcgtcag aatccccaga 120
gatgttcttg tttctggtta ttatttctgg ggtaatccaa ctcttgagaa gaagccagac 180
tcaccgggaa cttatgttga atggttccctc aaaggaaatg ttctatatgg atcatgggtt 240
gagcacatcc gtgcctgggt gtccatgcaa gaatgggaca acttcttggt actgtactat 300
gaag 304

```

```

<210> 60
<211> 293
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700302770H1

```

```

<400> 60
gtagccactc tascctagggg cgtgctgaga caagaccacc tcattcctct gctgcttttc 60
agacaggact gtccctgcga cccaccatga tccaggctgc actgttctt ggctgtatct 120
tactgtctct ggtgaccgcc ttctcatgga agactcagga tggctggcctg ccccatcaga 180
cagctggcac agaaactgag cctacacaac tgctctacag caagagtctt cctccgacct 240
ccagtaacctg tgggaaacct ctaagcatgg cgcacctgcc cctgtagtc etc 293

```

```

<210> 61
<211> 174
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No: 700303111H1

```

```

<400> 61
caacagcaaa cgtgcgacca actcttcggc tagtgaatct ctaagccgcy aagagtgttt 60
tgaagtagct ttaggtggaa gatgtcagaa agtaactcgg cagagggtag cgcagaagc 120
gaggagccag bgtctggctg taaagtcac gccacggccc taaaaacgca agat 174

```

```

<210> 62
<211> 273

```

<212> DNA  
 <213> Rattus norvegicus

<220>  
 <231> unsure  
 <222> 52, 139, 249  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700303390H1

<400> 52  
 ctcagatggc aggcattggca tgcattggat cttgttccct gagacaaagg cngatgcaga 60  
 gggcatgtga ataatcatg aggggcccac agcaggccag caggccatag ctgacctcat 120  
 tctggaagtg agagtggang agaccccagc tgggacagaa aaggtaaccac gctataaacc 180  
 atggcctaac cgaggggccag cagtggcagc ctccctgaaa gggacttcca gtccatccac 240  
 aggcaccgna gaaccagcaa gacatagcaa gcc 273

<210> 63  
 <211> 279  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700303722H1

<400> 63  
 gtgactctga gtgtttgagc aggttaacttc taacttttgc cctctatctgc aacagggtcca 60  
 aagggttccaa aggagctggc aggaacttca gacaagatcc actggcttca ggtgtgecta 120  
 gtcttgaggt tcagaaagac ggaggcagct gaattgtgtg ctgaaccaac aacatctagc 180  
 tacaagggga gccactcttc cccccagcga ctgtgactgt tctcaccagt ctgaatttcc 240  
 tgtttgtatt cacaagatg ctttttattt ttaacttct 279

<210> 64  
 <211> 275  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700306343H1

<400> 64  
 gagaaaggcc accacctagc taggtgaggt gtgccagcat agtccctggg gtctcactgt 60  
 ccccagccct gggacgtatg ttccgccatg caatcccttt cgtatcttc aagctgttcc 120  
 ttctttatct cagtgtatgg ctcttccatg agtggccctt tgagttgcca gtccaaagaa 180  
 ctcagcagtc cggcctgttg gaactcaagc tctctctctc ttctccagcc ctcacctctc 240  
 tgcctcctgt cactcaggt gttttacag gctga 275

<210> 65  
 <211> 294  
 <212> DNA

<213> Rattus norvegicus

<220>

<221> misc feature

<223> Incyte template ID No: 700306615H1

<400> 65

```

catccgtggg ctggctcag ccattcgct gtctctggag tatcacagaca caagctatga 60
ggacaagaag tacagcatgg gggatgctcc cgactatgac agaagccagt ggcctgagtga 120
gaagttcaaa ctgggctgg acttccccc ttctgacctac ttaattgatg ggtcacacaa 180
gatcaccag agcaatgcca tctgctgcta ccttggccgg aagcacaacc tttgtgggga 240
gacagaggag gagaggattc gtgtggagct ttgggagaac caggctatgg acac 294

```

<210> 66

<211> 283

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 2

<223> a or g or c or t, unknown, or other

<220>

<221> misc feature

<223> Incyte template ID No: 700330359H1

<400> 66

```

gngggccaat ggcccttgag taaaagggtg tcaatgagag tctaaggcc agagtaggaa 60
catgcggctt agagccagtg tcttgaccca gaacgtcac tcttgatag gtagatgagg 120
aggtgttgg gtgcccag gggtatccc gctgtgcttt cgccttagcc tttctgcaac 180
gcatggacat gaagccgtg gtggtcctgg gactgccggc ccggacggcc ccttcgggt 240
gtctctcctt ctgggaagct aaggcacagc ttgctcagag ctg 283

```

<210> 67

<211> 263

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc feature

<223> Incyte template ID No: 700368974H1

<400> 67

```

tccttagctg ggggoggggg gggcagtcg ctactatctg aaagaattta atgtgggagt 60
catgccttat acaacacagg aaattaatgt gtgatcta at gcgtgatcta tgaattatta 120
caatacagga atttaatgtg tgagccatgc ctccaaaaca tgtctagaat ttctggaatt 180
ggcgggaagt caacagggat tgcttatita acccttccaa tcaactcattg tgactagggc 240
acatggtctt ggccttgcta tga 263

```

<210> 68

<211> 269

<212> DNA

<213> Rattus norvegicus



<220>  
 <221> unsure  
 <222> 129, 133, 153, 163, 165  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700373118H1

<400> 68  
 gtttgaagca gggcgtaggg asaagcggga ttaagagga ctacettat tagctcttc 60  
 cctccagaat aggtgcaaat cctcccccatt gccatttcc tgcacactgg ggtaaggatg 120  
 tggcactgnc agnctgtcag cccactgact tttagctcttc agntngcagt ctgggcaaat 180  
 accagcgagc tcbgttgaa caagaccagg ccttcagagc atctgaacca ctgtggcctt 240  
 ctctcctcag ccttcactgt ggttttggc 269

<210> 69  
 <211> 288  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 159  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700375521H1

<400> 69  
 gcagccatgg atcgggggga ggaacctctg tcccgagagc cggcgctgga gaccgagagc 60  
 ctgcgattcc tgcaggtcac agtgggctcc ctgctggcca gctatggctg gtacatcctc 120  
 ttccagctgg tcbttctcta cattgtcacc cagaagctnt cctcgcgact gagggcttta 180  
 aggcagaggg agctggacca agctgaggct gttctggagc ctgatgttgt tgttaagcga 240  
 caagaggctt tagcagctgc tegtctgaga atgcgggaag atctgaat 288

<210> 70  
 <211> 280  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 16, 84, 96, 112-114, 118, 171, 226, 228, 236, 240, 267  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700461546H1

<400> 70  
 tgaacaaact tctganaact tagttgacag tgttttgagt caactgaaa aagcatgact 60  
 tttgggaatct ctgaatgact tggntctcag tattancaat ctattgaatt tnnctctnat 120  
 taaagtatgt agttttttag acctttttcc tgacagctatt atgtaatttt ntggcgctggg 180

tagatgggag tgcgcgttgc atgttaccat acagctgaca tctatctntt gctatctctn 240  
attatcttag tsgtttcatg ctgtgggctg taacataacc 280

<210> 71  
<211> 271  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 25, 64, 70, 82, 102, 155  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700480077H1

<400> 71  
ctgacctgac ccatgatgta agggncogta ggggagcctc accactgcaa aggcctgacta 60  
aggnctgttn ggctaaagggt cncctttgaag cccagtgtct anagtcacac cttctttgct 120  
ctgggcccag gaggcctact tcttcttttt ctcgnggaat cctgggaatct taaagataaa 180  
agaacctaga aagaaaatca aaccactttt ccttctgggg cagatggtas tatgggactg 240  
agaaccagca accctggggtc ttggagagga g 271

<210> 72  
<211> 210  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 187  
<223> a or g or c or t, unknown, or other

<220>  
<221> misc\_feature  
<223> Incyte template ID No: 700480949H1

<400> 72  
gggcagaggc ccagggaata agggaggctt ctaccaatga ttttgcttaa tggctgcttga 60  
cagagatatt gtatggttct ctgagagctc ccttgaaaac cttacctoca accacacaag 120  
ggttactccc agagagcgtc cgtggggcag caaggacaca ctcccatatt gccaaacata 180  
tcaagtnccc aaagattgga agaaaattcg 210

<210> 73  
<211> 256  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> unsure  
<222> 70  
<223> a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700483259H1

&lt;400&gt; 73

```

gtgaagctaca tggaaaacaa agcctaaggg gacaggctca agccgcagac agcagcaagt 60
aaagcgctctt cggccctgaa gcabggcagc tatcccttc agcggtctgc tcttggtac 120
ccatgactac tatcggtga agtagccct cgcagcccc gccagggct ggcccagggc 180
tctgtggtg acagacttc ccttcccagg acgtctgggc tctctgtca gcaacagctc 240
ggcgggaagt gcagag                                     256

```

&lt;210&gt; 74

&lt;211&gt; 259

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 219

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700483475H1

&lt;400&gt; 74

```

ctctgtact tggcaatgy ctggattccc accgtatca cggccttctt ccttgotacc 60
tcccaggccc aggttggtg gctacaacat gattatggc acctttctgt ctataagaaa 120
tccatattga accacattgt ccacaagttt gtcatgggc accttaagggt tgcctccgcc 180
aactggttga accatagaca ttccagcac catgogaano caacatcttc cacaaggacc 240
cgacataaa gagctgca                                     259

```

&lt;210&gt; 75

&lt;211&gt; 264

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700498995H1

&lt;400&gt; 75

```

gacagtagat gcccccaaa ctctagtaca tgatagtgtg ggggctgtgt ggggtctcta 60
ctctgtctgt tcattccag tgcagtttaa gggagcaggc gccactgcac tcttggctg 120
tgcctgagg gtgttgctg ctttatatag taacagtcaa ttaaggtttc ttccaggga 180
agaaaaggga tggttttgag gggctcagaa aataggatc agtgtgtaac ataacaggta 240
ggttgtcggc acatgctgat atcc                                     264

```

&lt;210&gt; 76

&lt;211&gt; 271

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

<221> unsure  
 <222> 218, 228, 255, 270  
 <223> a or g or c or t, unknown, or other

<230>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700504333H1

<400> 76  
 gtattatttg acacagacat ggacaaagcg atggagcgct atgtctctat gcccaaggaa 60  
 aaggctccag aaacacattcc cctctctctc attgcoctcc catcaagcaa ggatccaaac 120  
 tgggaggacc gattccacga ccgataccaca atgactgtgc tggtaaccac ggccctttgaa 180  
 tggttcgagg agtggcagga ggagcctaag ggcaagcnaa gtgttgcntt ggaaccccca 240  
 aaaaaacttc ccggnaaccc ttatatggga a 271

<210> 77  
 <211> 187  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 11-12, 17, 21, 24, 48, 66, 72, 96, 128, 135, 162, 166  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700505040H1

<400> 77  
 caacaattct nngtggncgt ncintctgga actgggcata atcagctnat gctgacatac 60  
 gctgntgga gngccgtggg gtgaaggteg ccgtnccct ggtgggtacc ttcattgtcag 120  
 cactaganat gctngtgtt tcccttactt tgatgcttgt gnatgna 187

<210> 78  
 <211> 267  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 5, 18, 26, 39, 90, 92, 122-123, 132, 137, 145, 152, 160, 168, 171  
 <223> 173-174, 186-187, 213  
 <223> a or g or c or t, unknown, or other

<230>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700505433H1

<400> 78  
 ggggncctct gtgaggcnet gatcncata gaggtgtga ttcagccagg ccacatgaag 60  
 ccccaagatg ggtggctctt cctgtatgan tngtaccaga tatatccatg gccggggaat 120  
 tnnagctggc anggtcncaa gggancacca gncagctttn tcaagaantc ntngggttc 180  
 cttggmctca caggaaaccta ttacctttca tgnngtctgg ggttctggat ttagggctct 240  
 tgggacagtc ccagttagaa gctctgg 267

<210> 79  
 <211> 267  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 14, 20, 22, 24, 31, 248, 253  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700510534H1

<400> 79  
 tgaagaggtta aggncttgcg tntnagaggc tctgtagtga gttctgtttg cctataaggg 60  
 aagtgggaaca accgagacac ncccacttct ttcgagtggtt aaggagcctg ggaggagcag 120  
 gcagcgcctt gctttgagca tgcacagggtg gggctgtccg ccgctgtggg gaaggcacc 180  
 tgcagcaggg ctctctgcc cactctctca ttgtagtgt gtccagatct cagaaacgca 240  
 gcttgaaacc agnttcaag gtaccag 267

<210> 80  
 <211> 291  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700513027H1

<400> 80  
 gggagggtttt aaaggccata ttgccaacct caaccgaagg ttccagggaac ccgagggaagt 60  
 gttaagtac aactcaccac ttcaagccac ccgaggctga agttgacgtt gcttttcaag 120  
 cctttttaca tacaactggc atttcagaaa attctcaaca ataattgtcty ccttcaggtt 180  
 taagtcatgy tgttttttag aattgacttg aatggaact atcaaaaagt gaatatatca 240  
 gctggtgac gagtgactga aacccccctg gtctgagggt gaccagttca g 291

<210> 81  
 <211> 273  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700528082H1

<400> 81  
 ataaaagtga aaactgggca agggcagggg gctggggctg aacagcttac tagataatgt 60  
 tctctaaaaa ttggtcttga aaacctgtt tgtgtattcg ttttatgagt gcttaaaaat 120  
 ggtgtgacca gggcatggc actgtcattg gaacagcaac atgcttgcctg gtcacattgg 180  
 aatgggggaa tgtgaagaa gctggacatc aggcctgcgg caccatttc ttgtatgaa 240  
 agtgttgtgt acaaaccccc cactaatcat ttt 273

<210> 82  
 <211> 268  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 68, 121, 165, 174, 182  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700520176H1

<400> 82  
 caaggccctc aggaatcag aggagccagc ctgacccctg gtctctggag tcttaaaaca 60  
 agtctgtntt tgcaggtagt cctagtggg tctgggggg aggetgccag gcttggcatg 120  
 ngacattagg caggaagcca ctctggatga ttgtgcacat gaganccctag tcaaggaggg 180  
 aagggtttta ggagaggact tgaattcaa gtgagaagcc agcagaggaa agggaaacca 240  
 gtccctcagaa tagaaggcta tactggct 268

<210> 83  
 <211> 289  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700534427H1

<400> 83  
 aaaacaggca agactttgga gaaagccagc cagggtatgat ggccactttg ccaccaacag 60  
 cccacacttc ccagcaacct gtaaacatag aggaogaaga tgggatccctg gatgagtatg 120  
 accagtacag cctggcccaa tcttatgttg tgggtggagg tgggaaaggc cgtaccaaga 180  
 gagaagctgc tgcacaacc aaccgcccc gacctgggtg gcattgagagg aagctgctga 240  
 ccaagttcca gaactctgaa aggaaaaagg cctggcgctg agacagagc 289

<210> 84  
 <211> 290  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 58, 157  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700535328H1

<400> 84  
 ggtctgacca agtgagaaag acagcagggg caccagggcc ccagacactc tggcgtantc 60  
 ccaaagaaag atggccacag cccagctccc tggtaaccaag ctgtcatccc taaaetctgc 120  
 tctgtgcccc tcttgggcag acgttaatca agcctngcc ctttctgatg ggcccccca 180

tcccggggaac actaaaaggt agtcttactg tcccccaccc tacaactgtt ttccataagtt 280  
atgcacaaat gcgaacagct gagacagaga tggagaagtt ctctgttttg 290

<210> 85

<211> 275

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 17

<223> a or g or c or t, unknown, or other

<230>

<231> misc\_feature

<232> Incyte template ID No: 700607163H1

<400> 85

cacagcccoct accagencac cctccataac tgcaccaaga ggaatctatcc aacacctccc 60  
tgagcaggag gagcctgaag actccaaggg aaagagctct gaggaacccct ttctgtgca 120  
gctggatcta accacaaacc cacaggggtga cacactggat gtctccttcc tctacctgga 180  
gcctgaggag aagaaactgg tggctctgcc ttctcctggg aaggaaacagc gctccctga 240  
gtgcccgggg cccgaaaagc aaagaacccc ctgat 275

<210> 86

<211> 285

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<222> Incyte template ID No: 700607235H1

<400> 86

ctgaagaccc accatgtctc tctgactac tgtactactt ctctgggggtt tcattctggg 60  
cccagcaact gacacagcct gtatattcaa ggaagcctcg gaaaacagtc ccttgcccag 120  
gcccgggctt cctgccaate cagtgcctcg gatcacact ggcctgaggc cattcctgct 180  
gtgccagggg acagtcgggg atgtactctt catgctgagg cgggaaggag atgatggctt 240  
cctggcgata gtccaacaga tgtttttctg gagggagctg gacc 285

<210> 87

<211> 260

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 246-247

<223> a or g or c or t, unknown, or other

<230>

<231> misc\_feature

<232> Incyte template ID No: 700607396H1

&lt;400&gt; 87

```

ggccaccaag atggcggggc ccagcgggacg gtgcgcgagg ttagtcagct gacttggtct 60
cggagctgtg ggcgcgcccc gcttctacct gtcccgagtg accagagctc agtgaccagt 120
ccttatagtc gaagcgaggg tttttactgc tgaggacctg gcccgctgg gaggccttgc 180
atggtaacag aacccgggggt agaggccata gggaaaaacc tagtggactc cagcgagccc 240
ctgcannccc gcttccgtgc                                260

```

&lt;210&gt; 88

&lt;211&gt; 181

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 27, 180

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700607505H1

&lt;400&gt; 88

```

caaaagaaaga aaacactcct ccgaggnccg cagcaagagg agagaaagat ctgcaggatg 60
accttcatta cagcaggtgt tatattttat cttttttgca cccgtttcta gtgaatgtat 120
cactaaggtc ttcaaaagaca tcagctttca aggaggtgca taagtactgt ttccacacaa 180
c                                181

```

&lt;210&gt; 89

&lt;211&gt; 280

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte template ID No: 700607713H1

&lt;400&gt; 89

```

aattgagggg taacatgtgc ttgttgaaat atatgttctt ttacaatact ttgtactaat 60
ttacgtggaa ttattatttg tttctcattg gagatattta ttacgctca atggccttc 120
aggaactcct gaateaggta ggaggcctag ggagattcca gatccttcag atggttcttg 180
ttgtcttcac cagtgttatt gtggtacctc atattataat agagaactta ctgcagccat 240
cccagtcct cgtgtctggg ttcttatcct cgacaatgtc                                280

```

&lt;210&gt; 90

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 6, 77, 128, 143, 146

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt;



<221> misc\_feature

<223> Incyte template ID No: 700607873H1

<400> 90

```
gaaganacac caaagctcac tcaactctca ggttctcaact gaacctactg catcggaagg 60
gaattcaact cccagangtc cttttttatg aagactgttg agacagcttt ccagaaacta 120
gaaccatntg gaagatagac ctngngtat tctgtgcgg attatcttga ttacgttaat 180
taattctgga tgggactagg cttaaagtgc atcatgattt tccattcaaa aggtgcacag 240
atgetacaaa tggctggggag aaatcct 267
```

<210> 91

<211> 258

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 11, 109

<223> a o r g o r c o r t, unknown, or other

<220>

<221> misc\_feature

<223> Incyte template ID No: 700607972H1

<400> 91

```
gaggctaaag ntgtgcctcc tggctctctc ggggcagtcg cccgcgcgc cagaactttc 60
gttgacctca ggtcccgct gctggcgaag gaaggcgagg gccactgong tctggacagc 120
gtccgaagga agcgagctct ctggaggcgc cgtagtga gaggagtcgg ttgtcacgtg 180
acccaagggt agaccatgga tttccacaag ccgtctgtct gcttctggga gtggggcaag 240
aatatggttt ggtggggg 258
```

<210> 92

<211> 276

<212> DNA

<213> Rattus norvegicus

<220>

<221> misc\_feature

<223> Incyte template ID No: 700608519H1

<400> 92

```
caaccacactg catctgcctt acatgcacac ttacaccato atctatttcc cctcccgagg 60
tgtggcctca ggtttttcaa gatgaaggga acaagccata gtcactcttg gggacccct 120
caacactggt gctggaggga tgtggccaag accaogttgg gggcaagacc gagacttggg 180
gcgggactac aattgtggtt ggtggggcca ggactgaact cttagcctcc atagccagct 240
acactgtctg taacttttcc tcttccctg tgcagg 276
```

<210> 93

<211> 298

<212> DNA

<213> Rattus norvegicus

<220>

<221> unsure

<222> 35  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700608661H1

<400> 93  
 cttagtccct gtactctgag ggtaaggctc atcgntcagg atctattgct gctgcttctc 60  
 cagctggctt ctgtagagac tgacagagac aactttgcag gataaggtag ctatcaagat 120  
 gctccatttg cgaagttcac agatgctgca gatgttggag agctccctaa ggaastacct 180  
 tcttgagtc ttaaaggctt atgggaactgt cttccacatg aaccaggag cccattcaa 240  
 gctcaaggct ctggtggaca agtggcctga tttatcacag tggttgtccg tctc 295

<210> 94  
 <211> 293  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 16, 179  
 <223> a or g or c or t, unknown, or other

<221>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700609074H1

<400> 94  
 ggcgtgaggt tggagnagag cgtcaggcgc ctccggggaga agtttcatgg aaaagtgtcc 60  
 cccaagaag cagggctctt tatgaggaag ttggcagcg accacactgg agtcggggcg 120  
 tctatctgtt acgggctcaa gcagaaagat ggacaggagc tgagcaacga tttggacgnc 180  
 caggaccac cagaggacat gaaggaggac caagatatcc aggcagttag cactctctgt 240  
 ttgcctctga cgaagccaa tcttcgaatg ttccaaagag cccagatga cct 293

<210> 95  
 <211> 288  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 66, 138, 152, 236, 242, 252-253, 279  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700609967H1

<400> 95  
 ctgcctcgca gcccagagcg cgcgcctctc cagctccgcg tccggcttcc ccaaccaggc 60  
 ttattntggc tcccgaccgg gtgcagacc ctagcccggc ctccggcccag ctccgcccc 120  
 tgcgtacttt taattggag gaggtggcgc angctccggg agggagaggg agcgatggct 180  
 cstaatcgac cgaaggtgt acaacatcag cgaattcagt tgcgcgccc cggggnggct 240  
 mcgggtcat ccccactag ctggtcagga tgcacggna tcttttgt 288

<210> 96  
 <211> 164  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700625315H1

<400> 96  
 gttagagcag ttacactgaa ccaaaagtga tgagtttgta cagacggtaa tccgtaccaa 60  
 gcacactcac tgtcctgato tgaacaccca gcaaggttca bgtcogtgcg aagtttgcag 120  
 cattgtgttc ttttgcattc tttttttact tttattaaag gttc 164

<210> 97  
 <211> 225  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 28, 59, 65, 73, 79, 84, 88  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700627069H1

<400> 97  
 cggggcggct gcagcaggcc accatgggag agcttcagga ggtgcagatc actagaggng 60  
 aagcnattgt tgnacaggtnt ggcnttantg gttgagtota tcaactccagt gggttccctg 120  
 tttgtcttgg catcatactc catcatcttc ctcaagcttt tctcctacog ggatgtcaat 180  
 ctgtgggtgac gccagcgaag ggtcaaggcc aaagctgtgt ctgca 225

<210> 98  
 <211> 265  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 62, 264  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700629293H1

<400> 98  
 atgaccttta actttttctaa aaatgtgaag ttttgtactt atatatatca gctaaagtac 60  
 tntagcattc tttagtgtac ttagtttgat gccactttta gtgtttttgt tgccttttgc 120  
 tgatttttat gaatgttcat ttttagactc cttgttgaaa tgggacagtt tegtcttttg 180  
 ataagccoga gaagaggatt cccttgggtg ttgacctctc ctgcattgat tgcccaagca 240  
 tctgaactgc aaccaaggcc ttnc 265

<210> 99  
 <211> 95  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No: 700643961H1

<400> 99  
 agtataacca ggggcaatct gaaagtgtgt ctctagccag ataagccact atgagcggta 60  
 agcatcagt ctaacgcaga aacttctga gcagg 95

<210> 100  
 <211> 307  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 8, 20, 21, 23, 47, 302  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700061635H1

<400> 100  
 ggggaaganac actggcttan nanttggttc tgaggggaagc tctctgngtg gatcacatatt 60  
 tataggattt ggggtgcaag aactgtgtga ctagtctcagt tcaattcagg gagcaggcag 120  
 aaaccagcag gtattagaag agatgttcta tatacagagt tctgaggcac tgcagattct 180  
 gaagaatttc ctaaggaagc acctccctga gtccttaaaag gttatgggac tgtctccaca 240  
 tgaaccaggg aaaccatttc aagctcaagg ctgtgtgtgga caagtggctg atttaatact 300  
 gntgtta 307

<210> 101  
 <211> 97  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 48, 90  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700062809H1

<400> 101  
 cccctgagagt ctccaaaatg tttcagttctg ttataagAAC cccattanct caataaagct 60  
 tgcactggct gactctctga cggggttccn gaatgtg 97

<210> 102  
 <211> 214  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700139104H1

<400> 102  
 gagatgttcc ctgtcatcga acagtatgga gacattttgg taaaatactt gaggaagag 60  
 aaaggcaaac ctgtccctgt gaaagaagtg ttgtgtgctt acagcatgga tgtgatcacc 120  
 agcacatcat ttggagtga tgttgattcc ctcaacaaac cgaaggatcc ttttgtggag 180  
 aaagccaga agctcttaag aattgatttt ttgt 214

<210> 103  
 <211> 265  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700139953H1

<400> 103  
 aggaagccct gcagagcacc agaggccctag ctaggaggac aacacagagg agtaatttgc 60  
 tgaacagacc gcagggatgg aactgotttc agctctcaca ctggaaacct gggctctctc 120  
 ggcagtctgc ctggtgctcc tctacggatt tgggacccgc acacatggac ttttcaagaa 180  
 acaggggatt cctgggccc aacctctgcc tttttttggc actgtgatga attactatat 240  
 gggtttatgg aaattcgatg tggag 265

<210> 104  
 <211> 306  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 2, 61, 68, 111, 139, 263, (296)...(298), 305  
 <221> unsure  
 <222>  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700289281H1

<400> 104  
 ancagatccg ggcactggag agggagctgc aagcaggagc agtcaagsgt gtggtcagaa 60  
 ncacagtnag tggaccagca gggccggctt ttctaccat ggcggcccaa ngctatggct 120  
 attacgcac tgtcatatnc acagccatgt cgggaggcta cagcctttac tacttcaacc 180  
 gcaaacctc ctcttttctc atgacctctt tggtagatga gatcgtctct gacaaggacg 240  
 atttgggggc tcatcaogag canccagtcg gcagcctaag catcagcaag tttgtannag 300  
 gggtna 306

<210> 105  
 <211> 183  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700303922H1

<400> 105  
 ggacgacattg atccttatgt atctctgccc ttgggaatg gacccaggaa ctgcattggc 60  
 atgagggttg ctctcatgaa tatgaaactc gctctcacta aagttctgca aaacttctcc 120  
 ttccagcctt gtaaggaaac acagatacct ctgaatttaa gcagacaagg actttctcaa 180  
 cca 183

<210> 106  
 <211> 290  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 56  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700305792H1

<400> 106  
 ctacacgtga atacatagct gogatccctg aactcagctc cctcatagtg aaacgncaac 60  
 gccagccctt cctgtacctg gacttcctgt attgcctcac tgcctatggg cggcgcttcc 120  
 gcaaggcctg cgaactggtg cacaacttca cagatgctgt catcaggagg agacgcagca 180  
 cctcaataac ccaggcggtt gatgaattcc taaggccag ggctaagact aaaactttag 240  
 actctattga tcttctcttg ctggccaagg atgagcatgg gaaggggctg 290

<210> 107  
 <211> 177  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700329969H1

<400> 107  
 gctatatcag gaggggaccc atgctgtgtc cttctgagat ctaacaggat taaccaatat 60  
 gtaaaactaga ggaagtgtt ggctgcact gggcaagccc tctaggactc catccaagaa 120  
 agaccagttg gtgttgcct agaggcaag aaaccataa ggagcttgca gtaaac 177

<210> 108  
 <211> 188  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 114, 116  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700368493H1

<400> 108  
 aaagcttccc aatctgtgtg ctcaattggg aggatggcat taggagccag ggcaggccat 60  
 gggtaacctta ctttccctcc tgggttatgc ccaggagaat ggagaaaaa aaangnttta 120  
 aagaaaaaat attttaaatt tgatgttggc ctctttccat tgtattgagt aaaagtgttc 180  
 aagttgtc 188

<210> 109  
 <211> 255  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 156, 157  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700376694H1

<400> 109  
 ttatgattat caattttaca tcaattaat attatctcaa acctccttaa gaaaatgagt 60  
 atggatgttc acagtatgtt tgatttttat ctacaagaat gaatctgatt cagaatgctt 120  
 ttcagctgac atacagagca ctaatactt taaggnaac cataggtctg aatctcttaa 180  
 gaattctcag tctctatggg atgtagggac gcattataaa tgcattaato cttatagtaa 240  
 atcctgtgcc tagga 255

<210> 110  
 <211> 264  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 16, 25, 54, 63, 68, 70, 80, 141, 154, 210, 275  
 <221> unsure  
 <222>  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700483803H1

<400> 110  
 gaggaacgcc gccgctctgc tcygnatcct acacccaatca ggaagctgct gtcnagccat 60  
 ggnngganen gagaagccan ctcaagagga tgaactggaa cccatggtaa catcaggggc 120

```

ctcagaagca gtgccaaggg ngctttcttg agancctcag aacatctctg atgtagatgc 180
cttcaacttg ctcttgga tgaactgan acgacggcgt gaggtcccaa ccttcacatg 240
actgtgaccc agctagtggc cgaggatggc agcanggtgt atgt 284

```

```

<210> 111
<211> 250
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No.: 700502844H1

```

```

<400> 111
ggcagacatt agcagattct ggatgccatc agtgaactgaa gccattaaag agactcggct 50
ggagttagcag ctaagaggac agagagacaa gggctacgag gcagcaatat aaacagatct 120
ggtgttgctg agatttgaga cgaaggcttc ccattggcttc ttctcacatc cgcagttcc 180
aggagaggga ctatgaacag gtctgggata tgttctccag gggantgaag gaacacatcc 240
ccactgcctt ccgcact 258

```

```

<210> 112
<211> 250
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No.: 700503415H1

```

```

<400> 112
gtagctttcc ccttttctg gcacagaagt ctgtccatct gcaagcgctt tggaacacag 50
actgcctgga gccaccttcc ttggggagac ctctctgctt cagctgtcgt cctgtgtcgt 120
cattcaactaa agctcctgac gtcagattaa gcaagcagtg atgggttaca tttagagcaa 180
gccgcagaga caaggcctgt tgcctgtttc cagataatga tgagttttaa ttaccactg 240
gtttgtatgg 250

```

```

<210> 113
<211> 279
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 40, 44, (53)...(55), 60, 69, 72, 74, 159, 234
<221> unsure
<222>
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature
<223> Incyte template ID No.: 700514914H1

```

```

<400> 113
ccactgcagc gccccccccc aaagatggaa agattctgcn ttancttcac atnncttan 60

```



```

aacatttgna cncnattett taaattttaga atctcccccga gagcctgttc ttcttttaac 120
accttttatg ctggagtagg atgatggctg agttctctta aaagagctta aatataaggt 180
cccaaacatg agatagatgc ctgcgcgcga ctctttccac aaactcgaga accnctttgt 240
gcacgcagct gccatggaag gaattcctgg ggcttctt 278

```

```

<210> 114
<211> 308
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<223> Incyte template ID No.: 700519169H2

```

```

<400> 114
ccccatcttt taagactttt cataatgatg ttaagaccag ggcagactat ctactggcca 60
tggactgcag ccaggagatg cctgcccct cctgggacct gcacaccacc tctctgggga 120
acttgacaaa ggtccctaag gctaaggag gttctcttc tactaggtec ctgactttga 180
ctctgtggct ctctaggaa cgtatgcaca ctctctctg ttgtaaccac aaagggcagt 240
agcacttagc atgtcatgtc ctgcccggg tgettgcctc ccaccacacc aggattctct 300
gggctggc 308

```

```

<210> 115
<211> 124
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 34, 53, 87, 109, 122
<223> a or g or c or t, unknown, or other

```

```

<220>
<221> misc_feature
<223> Incyte template ID No.: 700535905H1

```

```

<400> 115
cttacctctg gtggaaaaag taccatcctc aggnccagct ggtaagttt gtngetgaca 60
atcatccaga ccagctggcg ggtcatatgg ccgtgtctct tccctctcng gtggctgtac 120
tccc 124

```

```

<210> 116
<211> 262
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> unsure
<222> 148
<223> a or g or c or t, unknown, or other

```

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700533964H1

<400> 116  
 gggaaagtgt gttctgaggg cctgtgggg ccaaggggga ccagcctcac attccacacg 60  
 tgcgcacac cgtttggagc ctatttatt tgtatttatt tgaacagagt tatgtcctaa 120  
 ctatttttat agatttggtt aattaatncc ctgtcatttt caagttcatt ttttttattc 180  
 atatttatgt tcatgggtga ttgtacctcc tgtcaccagc tggtagggga gggagacaa 240  
 ggtagaaagt ggccacagag tg 262

<210> 117  
 <211> '267  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> unsure  
 <222> 36, 37, 223  
 <223> a or g or c or t, unknown, or other

<220>  
 <221> misc\_feature  
 <223> Incyte template ID No.: 700607844H1

<400> 117  
 ctogatogtc cagaccccac gggtcacc cgtcgnncca tcatcaggaa ctcaactgttc 60  
 gggagcgttg agacgtggcc ttggcaggtt ttaagcaccg ggggcaagg aagcgtctcc 120  
 tatgaggaaa gagcctgoga agggggaaag tttgctaact tggaaagtga agacaaaact 180  
 gtggatgagg ctctcgggga agcaatgcac aagatcatga agnctgggg ggcacccatg 240  
 acaaaggagt cggcatggga atgacag 267

**PCT**WC INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12Q 1/68</b>		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/12760</b> <b>(43) International Publication Date:</b> 9 March 2000 (09.03.00)
<b>(21) International Application Number:</b> PCT/US99/19768 <b>(22) International Filing Date:</b> 27 August 1999 (27.08.99) <b>(30) Priority Data:</b> 09/141,825 28 August 1998 (28.08.98) US 09/172,711 13 October 1998 (13.10.98) US 09/172,108 13 October 1998 (13.10.98) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications</b> US 09/141,825 (CIP) Filed on 28 August 1998 (28.08.98) US 09/172,711 (CIP) Filed on 13 October 1998 (13.10.98) US 09/172,108 (CIP) Filed on 13 October 1998 (13.10.98) <b>(71) Applicant (for all designated States except US):</b> INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CUNNINGHAM, Mary, Jane [US/US]; 1204 Manet Drive, Sunnyvale, CA 94087		<b>(74) Agents:</b> BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <b>(88) Date of publication of the international search report:</b> 3 August 2000 (03.08.00)	
<b>(54) Title:</b> TOXICOLOGICAL RESPONSE MARKERS <b>(57) Abstract</b> <p>The present invention relates to a composition comprising a plurality of nucleic acid molecules. The composition can be used as hybridizable array elements in a microarray. The present invention also relates to methods for screening compounds and therapeutics for metabolic responses indicative of a toxic compound.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

Interr Application No  
PCT, 99/19768

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL [Online] embl heidelberg AC#H33194, 30 September 1995 (1995-09-30) LEE H.N. ET AL.,: "EST108960 rat PC-12 cells" X:002130308 abstract	6
X	& LEE N.H. ET AL.,: "Comparative expressed-sequence-tag analysis of differential gene expression profiles in PC-12 cells before and after nerve growth factor treatment" PROC. NATL. ACAD. SCI. USA, vol. 92, - August 1995 (1995-08) pages 8303-8307, XP002047703 the whole document --- -/--	6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

14 February 2000

Date of mailing of the international search report

19. 05. 2000

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Müller, F

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/19768

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 13877 A (LYNX THERAPEUTICS INC ;MARTIN DAVID W (US)) 17 April 1997 (1997-04-17) see whole doc. esp. claims and p.4, line 12 ff. ---	1-5,7,8
Y	WO 95 20681 A (INCYTE PHARMA INC) 3 August 1995 (1995-08-03) see whole doc. esp. claims and p.40, line 3 ff. ---	1-5,7,8
Y	SCHENA M ET AL: "PARALLEL HUMAN GENOME ANALYSIS: MICROARRAY-BASED EXPRESSION MONITORING OF 1000 GENES" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 93, no. 20, 1 October 1996 (1996-10-01), pages 10614-10619, XP002912238 ISSN: 0027-8424 the whole document -----	1-5,7,8

# INTERNATIONAL SEARCH REPORT

Inte. App. No. JS 99/19768  
PC

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-8 (partially)

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

**1. Claims: 1-8 (partly)**

Nucleic acid sequence (seq id 1), fragments thereof, methods for detecting and diagnosing metabolic effects of compounds using microarrays comprising seq id 1 or any fragments thereof and methods for screening a library of molecules for specific binding to seq id 1.

**2. Claims: 1-8 (partly)**

Nucleic acid sequence (seq id 2), fragments thereof, methods for detecting and diagnosing metabolic effects of compounds using microarrays comprising seq id 2 or any fragments thereof and methods for screening a library of molecules for specific binding to seq id 2.

inventions 3-117

...ibidem for each nucleic acid sequence or a fragment thereof separately seq id 3-117



# INTERNATIONAL SEARCH REPORT

Automatic patent family members

International Application No  
PCT/JP 99/19768

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9713877 A	17-04-1997	AU 712929 B	18-11-1999
		AU 4277896 A	06-05-1996
		AU 6102096 A	30-12-1996
		AU 7717596 A	30-04-1997
		CN 1193357 A	16-09-1998
		CZ 9700866 A	17-09-1997
		CZ 9703926 A	17-06-1998
		EP 0793718 A	10-09-1997
		EP 0832287 A	01-04-1998
		EP 0931165 A	28-07-1999
		FI 971473 A	04-06-1997
		HU 9900910 A	28-07-1999
		JP 11507528 T	06-07-1999
		JP 10507357 T	21-07-1998
		NO 971644 A	02-06-1997
		NO 975744 A	05-02-1998
		PL 324000 A	27-04-1998
		WO 9641011 A	19-12-1996
WO 9520681 A	03-08-1995	US 5840484 A	24-11-1998
		AU 688465 B	12-03-1998
		AU 1694695 A	15-08-1995
		BG 100751 A	31-07-1997
		BR 9506657 A	16-09-1997
		CA 2182217 A	03-08-1995
		CN 1145098 A	12-03-1997
		CZ 9602189 A	14-05-1997
		EP 0748390 A	18-12-1996
		FI 962987 A	26-09-1996
		JP 9503921 T	22-04-1997
		LV 11696 B	20-08-1997
		NO 963151 A	27-09-1996
		PL 315687 A	25-11-1996
		HU 75550 A	28-05-1997

Form PCT/ISA/210 (patent family annex) (July 1992)